

The spacetime positive mass theorem in dimensions less than eight

Michael Eichmair

Lan-Hsuan Huang

Dan A. Lee

Richard Schoen

October 18, 2017

Abstract

We prove the spacetime positive mass theorem in dimensions less than eight. This theorem asserts that for any asymptotically flat initial data set that satisfies the dominant energy condition, the inequality $E \geq |P|$ holds, where (E, P) is the ADM energy-momentum vector. Previously, this theorem was only known for spin manifolds [38]. Our approach is a modification of the minimal hypersurface technique that was used by the last named author and S.-T. Yau to establish the time-symmetric case of this theorem [30, 27]. Instead of minimal hypersurfaces, we use marginally outer trapped hypersurfaces (MOTS) whose existence is guaranteed by earlier work of the first named author [14]. An important part of our proof is to introduce an appropriate substitute for the area functional that is used in the time-symmetric case to single out certain minimal hypersurfaces. We also establish a density theorem of independent interest and use it to reduce the general case of the spacetime positive mass theorem to the special case of initial data that has harmonic asymptotics and satisfies the strict dominant energy condition.

1 Introduction

The following theorem is the main result of this paper. The technical terms are defined in Section 2.

Theorem 1 (Spacetime positive mass theorem). *Let $3 \leq n < 8$ and let (M, g, k) be an n -dimensional asymptotically flat initial data set that satisfies the dominant energy condition. Then*

$$E \geq |P|,$$

M. Eichmair: University of Vienna; e-mail: michael.eichmair@univie.ac.at

L.-H. Huang: University of Connecticut; e-mail: lan-hsuan.huang@uconn.edu

D. A. Lee: Queens College and CUNY Graduate Center; e-mail: dan.lee@qc.cuny.edu

R. Schoen: Stanford University; e-mail: schoen@math.stanford.edu

where (E, P) is the ADM energy-momentum vector of (M, g, k) .

We briefly survey earlier results: The special case of Theorem 1 where $k \equiv 0$ is called the time-symmetric case, or sometimes the Riemannian case. It is of particular importance. In the time-symmetric case, we have that $P = 0$ and the dominant energy condition becomes the assumption that the scalar curvature of g is nonnegative. The last named author and S.-T. Yau proved the time-symmetric case in dimension three in two articles from 1979 and 1981 [30, 31]. In [33], they extended their proof of the time-symmetric case to dimensions less than 8, as explained in detail in [27]. In 1981, they considered the general case $k \neq 0$ in dimension three and succeeded in proving that $E \geq 0$ by solving Jang's equation [32]. Later, E. Witten discovered a completely different proof that $E \geq |P|$ in dimension three [38, 25]. Witten's technique easily generalizes to all higher dimensions, as long as the manifold is spin [5, 12]. In dimensions higher than 7, a complication arises in the Schoen-Yau argument due to possible singularities of minimal hypersurfaces. Two different strategies for handling this complication have been announced by J. Lohkamp in a preprint [23] from 2006 and by the last named author in 2009. The first named author has generalized the spacetime $E \geq 0$ theorem to dimensions less than 8 (without spin assumption) in [16].

For earlier history of this problem, we refer to the introduction of [30]. The $E \geq 0$ theorem is sometimes called the positive mass theorem in the literature. We prefer to refer to it more accurately as the positive energy theorem. We reserve the phrase positive mass theorem for the $E \geq |P|$ theorem. This result could also reasonably be called the future timelike energy-momentum theorem.

Our proof of Theorem 1 is self-contained rather than by reduction to a previously known case. In particular, it gives a new proof of the $E \geq 0$ theorem for non-time-symmetric data. It follows from the work of D. Christodoulou and N. Ó Murchadha [7] that the $E \geq 0$ theorem implies the $E \geq |P|$ theorem in the vacuum case via a boost of the initial data slice in its spacetime development. At the end of this paper, we explain how our methods from Section 6 allow for a generalization of this boost argument to arbitrary initial data satisfying the dominant energy condition. This provides an alternative proof of Theorem 1.

Our main theorem does not include a characterization of the equality case $E = |P|$. The natural conjecture states that if $E = |P|$ in Theorem 1, then $E = |P| = 0$ and (M, g) can be isometrically embedded into Minkowski space with second fundamental form k . Our proof of Theorem 1 is by contradiction, so the analysis of the equality case will require a substantial new idea. We note that the so-called equality case of the Riemannian positive mass theorem is derived from the nonnegativity of mass, but that its proof is unrelated to the proof of nonnegativity of mass. The

situation in the case of general data is more complicated. It provides an interesting direction for future research.

The desired rigidity statement described in the preceding paragraph is already known to hold for spin manifolds. Although Witten sketched the basic idea for proving rigidity of spin manifolds in his 1981 article [38], a complete, rigorous proof in all dimensions was not given until the work of P.T. Chruściel and D. Maerten in 2006 [10]. Their argument is based on R. Beig and P.T. Chruściel's 1996 proof in dimension three [6].

We briefly review the minimal hypersurface proof of the time-symmetric positive mass theorem in [30, 27]. The argument is by induction on the dimension $3 \leq n < 8$. It proceeds by contradiction. Suppose that there exists an asymptotically flat Riemannian manifold (M, g) with nonnegative scalar curvature and negative mass $E < 0$. By a density argument [31], one may assume that (M, g) has harmonic asymptotics and positive scalar curvature. The harmonic asymptotics and $E < 0$ assumptions imply that the coordinate planes $x^n = \pm\Lambda$ are barriers for minimal hypersurfaces for all sufficiently large Λ . Consider an $(n - 1)$ -dimensional vertical cylinder ∂C_ρ of large radius ρ in the asymptotically flat coordinate chart. For every $h \in [-\Lambda, \Lambda]$ there is an area-minimizing hypersurface $\Sigma_{\rho, h} \subset C_\rho$ with boundary equal to the height h sphere on ∂C_ρ . If $n < 8$, this area-minimizing hypersurface is smooth. Every such $\Sigma_{\rho, h}$ lies between the barrier planes $x^n = \pm\Lambda$. The area $\mathcal{H}^{n-1}(\Sigma_{\rho, h})$ is minimized over h by some $h_\rho \in (-\Lambda, \Lambda)$. The corresponding surface Σ_{ρ, h_ρ} has the property that

$$\frac{d^2}{dt^2} \Big|_{t=0} \mathcal{H}^{n-1}(\Phi_t(\Sigma_{\rho, h_\rho})) \geq 0 \quad (1)$$

for any variation Φ_t of Σ_{ρ, h_ρ} that is equal to vertical translation along ∂C_ρ . One can then extract a smooth subsequential limit Σ_∞ of Σ_{ρ, h_ρ} as $\rho \rightarrow \infty$. This Σ_∞ is itself an $(n - 1)$ -dimensional asymptotically flat manifold with energy equal to zero. Moreover, Σ_∞ is a stable minimal hypersurface of M . Owing to (1), Σ_∞ is stable with respect to variations that are (sufficiently close to) vertical translations outside a compact set. Using the well-established relationship between the stability of minimal hypersurfaces and scalar curvature, the stability of Σ_∞ allows one to construct a conformal factor that changes the metric on Σ_∞ to one with zero scalar curvature. The stability with respect to variations that are close to vertical translations is then used to show that the conformal factor must decrease the energy of Σ_∞ , thereby violating the time-symmetric positive mass theorem in dimension $n - 1$. For the base case of the induction, when $n = 3$, one can show that the stability of Σ_∞ and its asymptotics at infinity are incompatible with the Gauss-Bonnet Theorem. When $n = 3$, choosing a special height h_ρ turns out to be unnecessary.

Our approach to the spacetime positive mass theorem is essentially a generaliza-

tion of the proof described above. In particular, it does not use the time-symmetric positive mass theorem as an input, as was done in the Jang equation approach of [32]. The proof is again by contradiction. Let (M, g, k) be an n -dimensional asymptotically flat initial data set satisfying the dominant energy condition $\mu \geq |J|$ and such that $E < |P|$. By our density theorem from Section 6, we may assume that (M, g, k) has harmonic asymptotics and satisfies the strict dominant energy condition $\mu > |J|$. We may assume further that P points in the vertical direction $-\partial_n$ of the asymptotically flat coordinate chart. The harmonic asymptotics and $E < |P|$ assumptions imply the coordinate planes $x^n = \pm\Lambda$ are barriers for marginally outer trapped hypersurfaces (MOTS) for all sufficiently large Λ . Again, we consider an $(n - 1)$ -dimensional vertical cylinder ∂C_ρ of large radius ρ . Let $h \in [-\Lambda, \Lambda]$. The results from [14] guarantee the existence of a MOTS $\Sigma_{\rho,h}$ whose boundary is equal to the height h sphere on ∂C_ρ . This MOTS is smooth if $n < 8$. Moreover, $\Sigma_{\rho,h}$ lies between the planes $x^n = \pm\Lambda$ and is stable in the sense of MOTS with boundary [20]. Since MOTS are not known to arise from a variational principle, there is no canonical way of singling out a suitable height h_ρ as in the time-symmetric case. To overcome this, we introduce a new functional \mathcal{F} on hypersurfaces with boundary on ∂C_ρ such that for some $h_\rho \in (-\Lambda, \Lambda)$ we (roughly) have that

$$\left. \frac{d}{dh} \right|_{h=h_\rho} \mathcal{F}(\Sigma_{\rho,h}) \geq 0. \quad (2)$$

Inequality (2) in conjunction with the MOTS-stability of $\Sigma_{\rho,h}$ plays a role similar to that of (1) in the time-symmetric case. Note that the h_ρ selected in the time-symmetric case by minimization would satisfy (2) in our more general argument. As before, we extract a smooth subsequential limit Σ_∞ of Σ_{ρ,h_ρ} as $\rho \rightarrow \infty$. This Σ_∞ is itself an $(n - 1)$ -dimensional asymptotically flat manifold with energy equal to zero, and Σ_∞ is a stable MOTS in M . Using the relationship between stability of MOTS and scalar curvature established in [21], one can construct a conformal factor that changes the metric on Σ_∞ to one with zero scalar curvature. Finally, (2) plays the role of (1) in establishing that the conformal factor must decrease the energy of Σ_∞ , thereby violating the time-symmetric positive mass theorem in dimension $n - 1$.

As in the time-symmetric case, the delicate height-picking argument is not required when $n = 3$.

The structure of the paper is as follows. Section 2 sets up the basic definitions and recalls some useful background material. Section 3 establishes the existence of the MOTS needed for the proof. Section 4 completes the $n = 3$ case of Theorem 1. The basic $n = 3$ argument, which is explained in detail in Sections 3 and 4, was first sketched out in [28, Section 7.2]. Section 5 contains the parts of the proof that are specific to dimensions greater than three, including the height-picking procedure. In

the last section, we show that an initial data set which satisfies the dominant energy condition can be perturbed by a small amount to one with harmonic asymptotics that satisfies the strict dominant energy condition.

2 Definitions, notation, and basic facts

Definition 1. Let B be a closed ball in \mathbb{R}^n with center at the origin. For every $k \in \{0, 1, \dots\}$, $p \geq 1$, and $q \in \mathbb{R}$ we define the *weighted Sobolev space* $W_{-q}^{k,p}(\mathbb{R}^n \setminus B)$ as the collection of those $f \in W_{loc}^{k,p}(\mathbb{R}^n \setminus B)$ with

$$\|f\|_{W_{-q}^{k,p}(\mathbb{R}^n \setminus B)} := \left(\int_{\mathbb{R}^n \setminus B} \sum_{|\alpha| \leq k} ((\partial^\alpha f)(x) |x|^{\alpha+q})^p |x|^{-n} dx \right)^{\frac{1}{p}} < \infty.$$

We usually write L_{-q}^p instead of $W_{-q}^{0,p}$.

Suppose now that M is a C^k manifold such that there is a compact set $K \subset M$ and a diffeomorphism $M \setminus K \cong \mathbb{R}^n \setminus B$. The $W_{-q}^{k,p}$ norm on M is defined in a routine way by choosing an atlas for M that consists of the diffeomorphism $M \setminus K \cong \mathbb{R}^n \setminus B$ and finitely many precompact charts, and then summing the $W_{-q}^{k,p}(\mathbb{R}^n \setminus B)$ norm on the noncompact chart and the $W^{k,p}$ norms on the precompact charts. The resulting space $W_{-q}^{k,p}(M)$ and its topology only depend on the diffeomorphism $M \setminus K \cong \mathbb{R}^n \setminus B$. This definition can be extended to the tensor bundles of M simply by considering components with respect to these charts. We usually write $W_{-q}^{k,p}$ for $W_{-q}^{k,p}(M)$ when the context is clear.

Definition 2. Let B be a closed ball in \mathbb{R}^n with center at the origin. For every $k \in \{0, 1, \dots\}$, $\alpha \in (0, 1)$, and $q \in \mathbb{R}$ we define the *weighted Hölder space* $C_{-q}^{k,\alpha}(\mathbb{R}^n \setminus B)$ as the collection of those $f \in C_{loc}^{k,\alpha}(\mathbb{R}^n \setminus B)$ with

$$\|f\|_{C_{-q}^{k,\alpha}(\mathbb{R}^n \setminus B)} := \sum_{|\alpha| \leq k} \sup_x |x|^{\alpha+q} (\partial^\alpha f)(x) + \sum_{|\alpha|=k} [|x|^{\alpha+q} (\partial^\alpha f)(x)]_\alpha < \infty.$$

Suppose now that M is a C^k manifold such that there is a compact set $K \subset M$ and a diffeomorphism $M \setminus K \cong \mathbb{R}^n \setminus B$. The space $C_{-q}^{k,\alpha}(M)$ can then be defined just as we did for $W_{-q}^{k,p}(M)$ in the preceding definition.

Definition 3. Let $n \geq 3$. An *initial data set* is an n -dimensional manifold M equipped with a complete C_{loc}^2 Riemannian metric g and a C_{loc}^1 symmetric $(0, 2)$ -tensor k . On an initial data set, one can define the *mass density* μ and the *current density* J by

$$\begin{aligned} \mu &= \tfrac{1}{2} (R_g - |k|_g^2 + (\text{tr}_g k)^2) \\ J &= \text{div}_g k - d(\text{tr}_g k). \end{aligned}$$

We say that (M, g, k) satisfies the *dominant energy condition* if

$$\mu \geq |J|_g.$$

It is often convenient to consider the *momentum tensor*

$$\pi = k - (\text{tr}_g k)g.$$

It contains the same information as k since $k = \pi - \frac{1}{n-1}(\text{tr}_g \pi)g$.

Let

$$p > n, \quad q \in ((n-2)/2, n-2), \quad q_0 > 0, \quad \text{and} \quad \alpha \in (0, 1-n/p].$$

We say that an initial data set (M, g, k) is *asymptotically flat*² of type (p, q, q_0, α) if $g \in C_{loc}^{2,\alpha}(M)$, $k \in C_{loc}^{1,\alpha}(M)$, and if there is a compact set $K \subset M$ and a diffeomorphism $M \setminus K \cong \mathbb{R}^n \setminus B$ for some closed ball $B \subset \mathbb{R}^n$ such that

$$(g - \delta, k) \in W_{-q}^{2,p}(M) \times W_{-1-q}^{1,p}(M)$$

where δ is a smooth symmetric $(0, 2)$ -tensor that coincides with the Euclidean inner product on $M \setminus K \cong \mathbb{R}^n \setminus B$, and such that

$$\mu, J \in C_{-n-q_0}^{0,\alpha}(M).$$

If (M, g, k) is asymptotically flat, one can define the *ADM energy* E and the *ADM momentum* P as

$$E = \frac{1}{2(n-1)\omega_{n-1}} \lim_{r \rightarrow \infty} \int_{|x|=r} \sum_{i,j=1}^n (g_{ij,i} - g_{ii,j}) \nu_0^j d\mathcal{H}_0^{n-1}$$

$$P_i = \frac{1}{(n-1)\omega_{n-1}} \lim_{r \rightarrow \infty} \int_{|x|=r} \sum_{j=1}^n \pi_{ij} \nu_0^j d\mathcal{H}_0^{n-1} \quad i = 1, 2, \dots, n.$$

Here, the integrals are computed in the coordinate chart $M \setminus K \cong_x \mathbb{R}^n \setminus B$, $\nu_0^j = x^j/|x|$, \mathcal{H}_0^{n-1} is the $(n-1)$ -dimensional Euclidean Hausdorff measure, and ω_{n-1} is the volume of the standard unit sphere in \mathbb{R}^n .

Remark. Theorem 1 still holds if we allow multiple asymptotically flat ends in the definition of initial data sets. We simply use the large celestial spheres in the other ends as barriers in the proof of Lemma 6.

In order to carry out our main argument, we require better asymptotic behavior.

²There are several incompatible notions of *asymptotic flatness* in the literature.

Definition 4. Let $n \geq 3$ and let (M, g, k) be an n -dimensional asymptotically flat initial data set. We say that (M, g, k) has *harmonic asymptotics* if there exists a $C^{3,\alpha}$ diffeomorphism as in the definition of asymptotic flatness, as well as a $C_{2-n}^{2,\alpha}$ function u and a $C_{2-n}^{2,\alpha}$ vector field Y such that for $i, j = 1, 2, \dots, n$,

$$\begin{aligned} u(x) &= 1 + a|x|^{2-n} + O^{2,\alpha}(|x|^{1-n}) \\ Y_i(x) &= b_i|x|^{2-n} + O^{2,\alpha}(|x|^{1-n}) \\ g_{ij} &= u^{\frac{4}{n-2}}\delta_{ij} \\ \pi_{ij} &= u^{\frac{2}{n-2}}[(L_Y\delta)_{ij} - (\operatorname{div}_\delta Y)\delta_{ij}] \end{aligned}$$

where a, b_1, \dots, b_n are constants, δ_{ij} is the Euclidean metric, and L_Y is the Lie derivative. The expression $O^{2,\alpha}(|x|^{1-n})$ stands for a function in the weighted Hölder space $C_{1-n}^{2,\alpha}$.

Notation. Let (M, g, k) be an n -dimensional initial data set. Let Σ be a two-sided $C^{3,\alpha}$ hypersurface with boundary in M with unit normal ν . Let D denote the ambient covariant derivative. We define the second fundamental form B_Σ and shape operator S_Σ of Σ using the convention

$$B_\Sigma(X, Y) = \langle S_\Sigma(X), Y \rangle = \langle D_X\nu, Y \rangle$$

for vector fields X, Y tangent to Σ , where the angle brackets denote the inner product g . We define the mean curvature scalar H_Σ to be the trace of S_Σ . According to this convention, the mean curvature of a sphere in \mathbb{R}^n with respect to the outward pointing unit normal is positive. We also define the *expansion*

$$\theta_\Sigma^+ = H_\Sigma + \operatorname{tr}_\Sigma k$$

of Σ , where $\operatorname{tr}_\Sigma k$ denotes the trace over the tangent space of Σ . If θ_Σ^+ vanishes on all of Σ , we say that Σ is a *marginally outer trapped hypersurface*, or *MOTS* for short.

Note that the property of being a MOTS depends on the choice of normal. For a vector field X defined along Σ but not necessarily tangent to it, we let

$$\operatorname{div}_\Sigma X$$

be the function on Σ which at $x \in \Sigma$ equals $\sum_{i=1}^{n-1} \langle D_{e_i} X, e_i \rangle$ where e_1, \dots, e_{n-1} is an orthonormal basis of $T_x\Sigma$.

Notation. Given a vector field X defined along Σ , we can decompose X into its normal and tangential components

$$X = \varphi\nu + \hat{X}.$$

Throughout this paper, whenever there is a vector field with the variable name X on a hypersurface Σ , the function φ and the tangent field \hat{X} are defined this way. We use η to denote the outward pointing unit normal of $\partial\Sigma$ in Σ .

The expression for the linearization of the expansion stated in the following proposition generalizes the well-known formula for the variation of the mean curvature. See, for example, [21].

Proposition 2. *Let Σ be a two-sided hypersurface with boundary in an n -dimensional initial data set (M, g, k) , and let ν be a continuous unit normal field along Σ . Let $X \in \mathfrak{X}(M)$ be a C^2 vector field, and let Φ_t be the flow generated by X . We can compute the expansion $\theta_{\Sigma_t}^+$ of the push forward $\Sigma_t := \Phi_t(\Sigma)$ with respect to the unit normal that points in the direction of $\Phi_{t*}(\nu)$ and pull it back to a function on Σ using Φ_t . The derivative of this function in t at $t = 0$ is denoted by $D\theta^+|_{\Sigma}(X)$. We have that*

$$D\theta^+|_{\Sigma}(X) = -\Delta_{\Sigma}\varphi + 2\langle W_{\Sigma}, \nabla\varphi \rangle + (\operatorname{div}_{\Sigma} W_{\Sigma} - |W_{\Sigma}|^2 + Q_{\Sigma})\varphi + \nabla_{\hat{X}}\theta_{\Sigma}^+, \quad (3)$$

where

$$Q_{\Sigma} = \frac{1}{2}R_{\Sigma} - \mu - J(\nu) - \frac{1}{2}|k_{\Sigma} + B_{\Sigma}|^2.$$

Here, k_{Σ} denotes the restriction of k to vectors tangent to Σ , and W_{Σ} is the tangential vector field on Σ that is dual to the 1-form $k(\nu, \cdot)$ along Σ .

Notation. Throughout this paper, we will drop the Σ subscripts when the context is clear. Everything is computed with respect to the metric g unless noted otherwise. In particular, we use \mathcal{H} to denote the Hausdorff measures associated with g .

Definition 5. Let Σ be a MOTS in an initial data set (M, g, k) . We define the operator

$$L_{\Sigma}v := -\Delta_{\Sigma}v + 2\langle W_{\Sigma}, \nabla v \rangle + (\operatorname{div}_{\Sigma} W_{\Sigma} - |W_{\Sigma}|^2 + Q_{\Sigma})v, \quad (4)$$

where v is a function on Σ . Although this operator is not self-adjoint, the Krein-Rutman Theorem shows that there is a unique (Dirichlet) eigenvalue with least real part. It is called the *principal (Dirichlet) eigenvalue* of L_{Σ} . This eigenvalue is real. If Σ is connected, the corresponding eigenspace is one-dimensional and generated by a $C^{2,\alpha}$ *principal eigenfunction* that is positive on the interior of Σ [20, p. 3]. If the principal eigenvalue is nonnegative, we say that Σ is a *stable* MOTS. This concept of stability was introduced in [20], based on the analogous definition for closed MOTS in [2]. It is easy to see that this generalizes the notion of stability of minimal hypersurfaces with boundary.

Proposition 3. *Let Σ be a stable MOTS in an initial data set (M, g, k) . For every compactly supported C^1 function v on Σ that vanishes along $\partial\Sigma$, we have that*

$$\int_{\Sigma} (|\nabla v|^2 + Q_{\Sigma} v^2) d\mathcal{H}^{n-1} \geq 0. \quad (5)$$

This follows from an argument in [21], *cf.* the proof of Lemma 15 below.

We state the following geometric variant of the strong maximum principle for ordered hypersurfaces that are subsolutions and supersolutions of the same prescribed mean curvature equation. We refer to [26, Lemma 1], [3, Proposition 3.1], and [4, Proposition 2.4] for similar results and partial proofs. It is important to pay attention to the choice of normal here. A good example to have in mind is the following: Consider a sphere of radius 2 that is tangent to a sphere of radius 1 in Euclidean space. Either the smaller one is enclosed by the larger one or it lies outside of it. The (obvious) conclusion of the lemma in this simple example is that the larger sphere cannot lie inside the smaller one.

Proposition 4 (Strong maximum principle). *Let g be a C^2 Riemannian metric on $M = \bar{B}_1^{n-1}(0) \times (-2, 2) \subset \mathbb{R}^n$. Let F be a C^1 function on the unit sphere bundle of M and let $u_1, u_2 \in C^2(\bar{B}_1^{n-1}(0))$ be such that $-1 \leq u_1(x') \leq u_2(x') \leq 1$ for all $x' \in \bar{B}_1^{n-1}(0)$. Assume that the hypersurfaces with boundary $\Sigma_i = \text{graph}(u_i) \subset M$ are such that $H_{\Sigma_1}(x) \leq F(x, \nu_{\Sigma_1}(x))$ for all $x \in \Sigma_1$ and $H_{\Sigma_2}(x) \geq F(x, \nu_{\Sigma_2}(x))$ for all $x \in \Sigma_2$ where the mean curvatures are computed using the upward pointing unit normals. If Σ_1 and Σ_2 intersect at an interior point or are tangent to each other at a boundary point, then they must be equal.*

Let g_1, g_2 be two metrics on an n -dimensional manifold M that are related by

$$g_2 = u^{\frac{4}{n-2}} g_1.$$

The scalar curvatures of these metrics are related by

$$R_2 = u^{-\frac{n+2}{n-2}} \left(R_1 u - \frac{4(n-1)}{n-2} \Delta_1 u \right). \quad (6)$$

Let $\Sigma \subset M$ be a two-sided hypersurface. If ν_1 is a unit normal with respect to g_1 , then $\nu_2 = u^{\frac{-2}{n-2}} \nu_1$ is a unit normal with respect to g_2 . The corresponding mean curvatures are related by

$$H_2 = u^{\frac{-2}{n-2}} \left(H_1 + \frac{2(n-1)}{n-2} u^{-1} \nabla_{\nu_1} u \right). \quad (7)$$

If (M, g_1) is an asymptotically flat n -dimensional Riemannian manifold and if $u = 1 + a|x|^{2-n} + O^{2,\alpha}(|x|^{1-n})$ is $C^{2,\alpha}$, then (M, g_2) is also asymptotically flat. The energies

of (M, g_1) and (M, g_2) are related by the formula

$$E_2 = E_1 - \frac{2}{(n-2)\omega_{n-1}} \lim_{r \rightarrow \infty} \int_{|x|=r} u \nabla_{\nu_1} u \, d\mathcal{H}_1^{n-1} \quad (8)$$

$$= E_1 + 2a. \quad (9)$$

The proof of the positive mass theorem in the time-symmetric case uses regularity and compactness properties of area minimizing hypersurfaces. By contrast, MOTS are not known to obey a useful variational principle. We will use the theory of almost minimizing currents as a viable substitute in our proof of Theorem 1.

Definition 6 ([13]). Let (M, g) be a complete n -dimensional Riemannian manifold and let T be an integral k -current in M . Let $U \subset M$ be an open set such that $\text{spt}(\partial T) \cap U = \emptyset$. Then T is λ -minimizing in U if for every integral $(k+1)$ -current X with support in U we have that

$$\mathbf{M}_U(T) \leq \mathbf{M}_U(T + \partial X) + \lambda \mathbf{M}_U(X).$$

Here, \mathbf{M}_U denotes the mass of a current in U .

This particular almost minimizing property was introduced and studied systematically by F. Duzaar and K. Steffen in [13]. In [14], the first named author of the present article observed that the λ -minimizing property is a natural feature of the MOTS that arise in the existence theory of the Plateau problem developed in [14], despite the absence of a useful variational principle. The properties of λ -minimizing currents that we use in the proof of Theorem 1 below are summarized in [14, Appendix A].

3 Construction of MOTS

Our proof of Theorem 1 will be by induction on dimension and contradiction. Let $3 \leq n < 8$, and suppose there exists an n -dimensional asymptotically flat initial data set (M, g, k) of type (p, q, q_0, α) satisfying the dominant energy condition, but $E < |P|$. For the case $n = 3$, we will obtain a contradiction to the Gauss-Bonnet Theorem in Section 4, and for $3 < n < 8$, we will obtain a contradiction to the time-symmetric case of Theorem 1 in dimension $n - 1$. By the density theorem (Theorem 18), we can assume without loss of generality, that (g, k) has harmonic asymptotics and satisfies the strict dominant energy condition $\mu > |J|_g$. Specifically, we can choose asymptotically flat coordinates (x^1, \dots, x^n) on $M \setminus K \cong_x \mathbb{R}^n \setminus B$ for some closed ball B such that on $\mathbb{R}^n \setminus B$, we have, for $i, j = 1, 2, \dots, n$,

$$g_{ij} = u^{\frac{4}{n-2}} \delta_{ij}$$

$$\pi_{ij} = u^{\frac{2}{n-2}} [(L_Y \delta)_{ij} - (\text{div}_{\delta} Y) \delta_{ij}]$$

for some $u, Y \in C_{2-n}^{2,\alpha}$ satisfying

$$\begin{aligned} u(x) &= 1 + a|x|^{2-n} + O^{2,\alpha}(|x|^{1-n}) \\ Y_i(x) &= b_i|x|^{2-n} + O^{2,\alpha}(|x|^{1-n}). \end{aligned}$$

Without loss of generality, we assume that $P = (0, \dots, 0, -|P|)$.

3.1 Existence of horizontal barriers

Lemma 5. *Let (M, g, k) be as described above. Then, for sufficiently large Λ , we have that $\theta_{\{x^n=\Lambda\}}^+ > 0$ and $\theta_{\{x^n=-\Lambda\}}^+ < 0$ where the expansion is computed with respect to the upward pointing unit normal.*

Proof. It follows from (9) that the $|x|^{2-n}$ coefficient of the function u is just $a = \frac{E}{2}$. We claim that the $|x|^{2-n}$ coefficient of Y_i is $b_i = -\frac{n-1}{n-2}P_i$, i.e. that $b_n = \frac{n-1}{n-2}|P|$ and $b_i = 0$ for $i < n$. To see this, note that

$$\begin{aligned} (n-1)\omega_{n-1}P_i &= \lim_{r \rightarrow \infty} \int_{|x|=r} \pi_{ij}\nu_0^j d\mathcal{H}_0^{n-1} \\ &= \lim_{r \rightarrow \infty} \int_{|x|=r} (Y_{i,j} + Y_{j,i} - (\operatorname{div}_\delta Y)\delta_{ij}) \nu_0^j d\mathcal{H}_0^{n-1} \\ &= \lim_{r \rightarrow \infty} \int_{|x|=r} (2-n)|x|^{1-n} [b_i(\nu_0)_j + b_j(\nu_0)_i - b_k\nu_0^k\delta_{ij} + O(|x|^{-1})] \nu_0^j d\mathcal{H}_0^{n-1} \\ &= -(n-2)b_i\omega_{n-1}. \end{aligned}$$

We claim that

$$\theta_{\{x^n=\Lambda\}}^+ = H_{\{x^n=\Lambda\}} + \operatorname{tr}_{\{x^n=\Lambda\}}(k) = (n-1)(|P| - E)\Lambda|x|^{-n} + O(|x|^{-n}). \quad (10)$$

To see this, we use harmonic asymptotics and formula (7) to compute

$$\begin{aligned} H_{\{x^n=\Lambda\}} &= \frac{2(n-1)}{n-2}u^{\frac{-2}{n-2}-1}\partial_n u \\ &= \frac{2(n-1)}{n-2}u^{\frac{-2}{n-2}-1}[(2-n)a|x|^{-n}x^n + O(|x|^{-n})] \\ &= -2(n-1)a|x|^{-n}x^n + O(|x|^{-n}) \\ &= -(n-1)E|x|^{-n}\Lambda + O(|x|^{-n}). \end{aligned}$$

To compute $\operatorname{tr}_{\{x^n=\Lambda\}}(k)$, first note that

$$k_{ij} = u^{\frac{2}{n-2}} [(L_Y\delta)_{ij} - \frac{1}{n-1}(\operatorname{div}_\delta Y)\delta_{ij}].$$

So we have

$$\begin{aligned}
\text{tr}_{\{x^n = \Lambda\}}(k) &= \sum_{i,j=1}^{n-1} g^{ij} k_{ij} \\
&= \sum_{i,j=1}^{n-1} u^{\frac{-2}{n-2}} \delta^{ij} [Y_{i,j} + Y_{j,i} - \frac{1}{n-1} (\text{div}_\delta Y) \delta_{ij}] \\
&= \sum_{i,j=1}^{n-1} u^{\frac{-2}{n-2}} \delta^{ij} [\frac{-1}{n-1} Y_{n,n} \delta_{ij} + O(|x|^{-n})] \text{ since } Y_i = O^1(|x|^{1-n}) \text{ for } i < n \\
&= -Y_{n,n} + O(|x|^{-n}) \\
&= (n-2)b_n |x|^{-n} x^n + O(|x|^{-n}) \\
&= (n-1)|P||x|^{-n}\Lambda + O(|x|^{-n}),
\end{aligned}$$

completing the proof of (10). Note that (10) shows that for large enough Λ one has $\theta_{\{x^n = \Lambda\}}^+ > 0$. The proof that $\theta_{\{x^n = -\Lambda\}}^+ < 0$ is similar. \square

3.2 Existence of MOTS

Notation. We now fix Λ large enough so that Lemma 5 applies. For large ρ , we define $C_\rho := K \cup x^{-1}\{(x^1, \dots, x^{n-1}, x^n) \mid (x^1)^2 + \dots + (x^{n-1})^2 < \rho^2\}$ to be the region horizontally bounded by a vertical coordinate cylinder of radius ρ , and we define $C_{\rho,h}$ to be the part of C_ρ lying between the planes $x^n = \pm h$. Define $\Gamma_{\rho,h} := \partial C_\rho \cap \{x^n = h\}$.

Lemma 6. *Let (M, g, k) and Λ be as described above. For every sufficiently large ρ and all $h \in [-\Lambda, \Lambda]$ there exists a stable $C^{3,\alpha}$ MOTS $\Sigma_{\rho,h}$ in $C_{\rho,\Lambda}$ whose boundary equals $\Gamma_{\rho,h}$ and which meets ∂C_ρ transversely. Every $\Sigma_{\rho,h}$ is a λ -minimizing boundary in $C_{\rho,2\Lambda}$ where λ depends only on $|k|_{C^0}$. Moreover, we can choose $\{\Sigma_{\rho,h}\}_{|h| \leq \Lambda}$ so that Σ_{ρ,h_1} lies strictly below Σ_{ρ,h_2} as a boundary in $C_{\rho,\Lambda}$ if $-\Lambda \leq h_1 < h_2 \leq \Lambda$.*

Remark. The regularity of the MOTS $\Sigma_{\rho,h}$ is the only place in the proof of Theorem 1 where the assumption $n < 8$ is used. For $n > 8$, the lemma still holds except that the λ -minimizing boundaries $\Sigma_{\rho,h}$ are only regular away from a thin singular set.

Proof. First observe that by the decay conditions on g and k , the coordinate cylinder ∂C_ρ has $\theta^+ > 0$ with respect to the outward normal and $\theta^- < 0$ with respect to the inward normal.

We would like to solve the Plateau problem for MOTS with boundary $\Gamma_{\rho,h}$ for each $h \in [-\Lambda, \Lambda]$. Note that $\Gamma_{\rho,h}$ divides $\partial C_{\rho,2\Lambda}$ into a top piece and a bottom piece. According to [15], the MOTS Plateau problem is solvable if the top piece has $\theta^+ > 0$ with respect to the outward normal of $\partial C_{\rho,2\Lambda}$ and the bottom piece has $\theta^- < 0$ with




Figure 1: The expansion θ^+ is computed with respect to the indicated unit normals.

respect to the inward normal of $\partial C_{\rho,2\Lambda}$. By the observation above and Lemma 5, the two pieces of $\partial C_{\rho,2\Lambda}$ satisfy the desired trapping conditions, with the exception of the corners where ∂C_ρ intersects $\{x^n = \pm 2\Lambda\}$. See Figure 1. Intuitively, the corners do not cause a problem because the (singular) distributional mean curvature there has a favorable sign. Therefore one could “round-off” the corners as in Figure 1. Alternatively, we observe that the proof in [14] can easily accommodate the corners: Just as in that proof, we use the trapping of the cylindrical and horizontal pieces of $\partial C_{\rho,2\Lambda}$ to construct barriers that have appropriate blow-up behavior for Jang’s equation, and then combine these barriers as in [15, Lemma 4.1]. Specifically, there exists a $C^{3,\alpha}$ family of MOTS $\{\Sigma_{\rho,h}\}_{|h|\leq\Lambda}$ with $\partial\Sigma_{\rho,h} = \Gamma_{\rho,h}$ such that each $\Sigma_{\rho,h}$ is a λ -minimizing boundary in $C_{\rho,2\Lambda}$ where $\lambda = \lambda(|k|_{C^0})$.

Moreover, it can be seen from the construction in [14] that the regions $\Omega_{\rho,h} \subset C_{\rho,2\Lambda}$ bounded by the $\Sigma_{\rho,h}$ ’s can be taken to be ordered, so that $\Omega_{\rho,h_1} \subset \Omega_{\rho,h_2}$ whenever $-\Lambda \leq h_1 \leq h_2 \leq \Lambda$. To see this, note that the supersolutions $\bar{u}_{t,\rho,h}$ used in the construction of solutions to the regularized Jang’s equation that lead to blow up in [14, Lemma 4.1, bottom of p. 570] can be taken to be pointwise decreasing in the parameter h , so that the corresponding Perron solutions $\bar{u}_{t,\rho,h}^P$ are decreasing in h and hence their epigraphs are increasing. It has been shown in [17] that we may assume further that $\Sigma_{\rho,h}$ is stable in the sense of MOTS.

Standard barriers for $\Sigma_{\rho,h}$ can be constructed from the trapped boundary ∂C_ρ by slight (C^2)-inward perturbation above respectively below its boundary $\Gamma_{\rho,h}$, locally uniformly in (ρ, h) , so that the angle (in the underlying Euclidean coordinate system) at which $\Sigma_{\rho,h}$ meets ∂C_ρ is bounded away from 0. For the special case of MOTS this inwards bending is explained in some detail in [14, Section 3]. Together with

the λ -minimizing property and Allard's boundary regularity theorem [1] it follows that near ∂C_ρ the hypersurface $\Sigma_{\rho,h}$ can be written as a vertical $C^{1,\alpha}$ graph above $\{x^n = h\}$. In particular, we see that $\Sigma_{\rho,h}$ intersects C_ρ transversely. Higher regularity of the defining function—which solves a prescribed mean curvature equation—then follows from Schauder theory in a standard way.

Finally, since each horizontal plane above $x^n = \Lambda$ has $\theta^+ > 0$ and each horizontal plane below $x^n = -\Lambda$ has $\theta^+ < 0$ by Lemma 5 it follows from Lemma 4 that each $\Sigma_{\rho,h}$ actually lies in $C_{\rho,\Lambda}$. \square

Remark. We are grateful to Brian White for helping us clarify the following issue: When M is an n -dimensional $C^{1,\alpha}$ manifold and g is a complete C^α Riemannian metric on it, then the standard interior Allard-type $C^{1,\alpha}$ regularity of (almost) minimizing boundaries away from a set of Hausdorff dimension at most 8 holds. This was shown by J. Taylor in [35] (this part of the discussion in her paper applies to n -dimensional manifolds). When the manifold is C^2 and the metric is Lipschitz, then this follows also from the work of R. Schoen and L. Simon [29] (for almost minimizers this was pointed out by B. White in [37, p. 498]). When the manifold is C^4 and the metric C^3 so that the Nash embedding theorem provides an isometric embedding of (M, g) into a high dimensional Euclidean space, then this also follows directly upon applying the Euclidean regularity theory as in [34]. In the preceding lemma, note that once we know that our surfaces are $C^{1,\alpha}$, we can then apply Schauder theory to the MOTS equation to obtain $C^{3,\alpha}$ regularity. The boundary regularity follows more easily because the metric is conformal to the Euclidean metric there.

3.3 Convergence of MOTS

Although there is no reason to expect the family $\{\Sigma_{\rho,h}\}_{|h|\leq\Lambda}$ to form a $C^{3,\alpha}$ foliation (even when there are no topological obstructions), we can still prove a partial regularity result. Its proof is similar to the proof of regularity of the outermost MOTS (established in [3] for $n = 3$ and then in [15] for $3 \leq n < 7$) but simpler, because the two-sided λ -minimizing property ensures embeddedness.

Lemma 7. *Let $\{\Sigma_{\rho,h}\}_{|h|\leq\Lambda}$ be as in Lemma 6. For each $h_0 \in (-\Lambda, \Lambda]$, the upper envelope of $\{\Sigma_{\rho,h}\}_{h < h_0}$ is a $C^{3,\alpha}$ MOTS with boundary Γ_{ρ,h_0} which we denote by $\underline{\Sigma}_{\rho,h_0}$. By convention we define $\underline{\Sigma}_{\rho,-\Lambda} := \Sigma_{\rho,-\Lambda}$. Moreover, $\lim_{h \nearrow h_0} \Sigma_{\rho,h} = \underline{\Sigma}_{\rho,h_0}$ in the $C^{3,\alpha}$ topology. We define $\overline{\Sigma}_{\rho,h_0}$ as the lower envelope of $\{\Sigma_{\rho,h}\}_{h > h_0}$ for $h_0 \in [-\Lambda, \Lambda]$ and $\overline{\Sigma}_{\rho,\Lambda} := \Sigma_{\rho,\Lambda}$ and note that analogous statements hold for these hypersurfaces.*

Proof. Fix $h_0 \in (\Lambda, \Lambda]$. Let $-\Lambda \leq h_i \nearrow h_0$ as $i \rightarrow \infty$ and pass the λ -minimizing boundaries Σ_{ρ,h_i} to a subsequential limit $\underline{\Sigma}_{\rho,h_0}$. Current convergence is automatic

from the mass bounds, varifold convergence follows because there is no mass loss in limits of λ -minimizing currents, and $C^{3,\alpha}$ convergence follows from Allard's interior and boundary regularity theorems. Note that since the Σ_{ρ,h_i} are increasing, this limit is independent of the choice of subsequence, and hence it is really a limit of the original sequence h_i . The limit does not depend on the choice of sequence h_i for the same reason. \square

Definition 7. We say that $h_0 \in [-\Lambda, \Lambda]$ is a *jump height* if $\overline{\Sigma}_{\rho,h_0}$ does not equal $\underline{\Sigma}_{\rho,h_0}$.

By the previous lemma, $\Sigma_{\rho,h}$ converges to Σ_{ρ,h_0} in $C^{3,\alpha}$ as $h \rightarrow h_0$ precisely when h_0 is *not* a jump height. It follows from the preceding lemma and Lemma 4 that h_0 is *not* a jump height if and only if the map $h \rightarrow \mathcal{L}^n(\Omega_{\rho,h})$ is continuous at h_0 , where $\mathcal{L}^n(\Omega_{\rho,h})$ is the volume of the enclosed region $\Omega_{\rho,h}$ defined in the proof of Lemma 6. Note that this implies that there are at most countably many jump times $h_0 \in [-\Lambda, \Lambda]$.

We also observe that the MOTS constructed in Lemma 6 may be used to construct complete MOTS with good asymptotics.

Lemma 8. *For any choice of $\rho_j \rightarrow \infty$ and $h_j \in [-\Lambda, \Lambda]$, there exists a subsequence of Σ_{ρ_j, h_j} that converges in $C^{3,\alpha}$ on compact subsets of M to a complete $C^{3,\alpha}$ properly embedded MOTS Σ_∞ . Moreover, there exists a constant $c \in [-\Lambda, \Lambda]$ such that outside a large compact subset of M , Σ_∞ can be written as the Euclidean graph $\{x^n = f(x')\}$ of some $C^{3,\alpha}$ function $f(x') = c + O^{3,\alpha}(|x'|^{3-n})$ in the $(x^1, \dots, x^{n-1}, x^n) = (x', x^n)$ coordinate system.*

Proof. Existence of a subsequential limit Σ_∞ and $C^{3,\alpha}$ convergence follow as in the proof of Lemma 7. Note that the limit Σ_∞ is again λ -minimizing. Since each Σ_{ρ,h_ρ} lies between the horizontal planes $x^n = \pm\Lambda$, so does Σ_∞ . The estimates below take part in the complement of a large ball B in M where we have harmonic asymptotics for the metric so that $g_{ij} = u^{\frac{4}{n-2}} \delta_{ij}$ where $u = 1 + O^{2,\alpha}(|x|^{2-n})$. It is not difficult to see from the corresponding property of Σ_{ρ_j} that the vertical projection of Σ_∞ onto the plane $\{x_n = 0\} \cap (M \setminus B)$ is surjective. Note that (7) implies that the Euclidean mean curvature of Σ_∞ is $O(|x|^{1-n})$. The λ -minimizing property of Σ_∞ gives rise to an explicit estimate of the form $O(|x'|^{-1})$ for the Euclidean area excess of Σ_∞ in large Euclidean balls centered at points $(x', x^n) \in \Sigma_\infty$ and of radius $|x'|/2$. Together with the Allard regularity theorem (the version in [34, Theorem 24.2] is particularly convenient here), this estimate implies that outside some large compact set, Σ_∞ is the graph of a function $f(x')$ such that $|f(x')| \leq \Lambda$ and $f(x') = O^{1,\gamma}(1)$ for some $\gamma \in (0, 1)$.

Since Σ_∞ is a MOTS we have that $H_{\Sigma_\infty} = -\text{tr}_{\Sigma_\infty}(k)$. As in [27, p. 32], this translates into a prescribed (Euclidean) mean curvature equation for f via a conformal change (7). The initial estimate $f = O^{1,\gamma}(1)$, together with a computation of $\text{tr}_{\Sigma_\infty} k$ as in the proof of Lemma 5, shows that the Euclidean mean curvature of f is $O^\gamma(|x'|^{1-n-\gamma})$ for some $\gamma \in (0, 1)$. Standard asymptotic analysis as in [24, 26] shows that there exists a constant $c \in [-\Lambda, \Lambda]$ such that $f(x') = c + O^{2,\gamma}(|x'|^{3-n})$. (Note that there is no logarithm term when $n - 1 = 2$ because f is bounded.) Repeating the above analysis with this information shows that $f(x') = c + O^{3,\alpha}(|x'|^{3-n})$, as asserted. \square

Corollary 9. *When $n > 3$, the hypersurface $\Sigma_\infty^{n-1} \subset M$ in Lemma 8 is asymptotically flat and has zero energy with respect to the induced metric g_∞ .*

The following lemma is a simple consequence of Proposition 3 and the fact that Σ_∞ is a limit of stable MOTS.

Lemma 10. *Let Σ_∞ be a complete MOTS whose existence is established by Lemma 8. For any $v \in W_{\frac{3-n}{2}}^{1,2}(\Sigma_\infty)$, we have*

$$\int_{\Sigma_\infty} (|\nabla v|^2 + Q_{\Sigma_\infty} v^2) d\mathcal{H}^{n-1} \geq 0.$$

We omit the proof because it is strictly simpler than that of Lemma 17 in Section 5.3. Specifically, the proof of Lemma 17 becomes a proof of Lemma 10 by simply replacing Z by an arbitrary compactly supported vector field on M and replacing the use of Lemma 15 by Proposition 3.

4 The case $n = 3$

We consider the base case $n = 3$ of our inductive proof. After the preparation of the previous section, the rest of the proof of the $n = 3$ case is essentially the same as for the time-symmetric case in [30], where minimal surfaces are replaced by MOTS.

By Lemma 8, we can extract a subsequential limit Σ_∞ of $\Sigma_{\rho,0}$ as $\rho \rightarrow \infty$. Let Σ'_∞ be the noncompact component of Σ_∞ . Lemma 10 implies that

$$\int_{\Sigma'_\infty} (|\nabla v|^2 + Q_{\Sigma'_\infty} v^2) d\mathcal{H}^2 \geq 0 \tag{11}$$

for every $v \in W_{\frac{3-n}{2}}^{1,2}(\Sigma_\infty)$. Noting that Σ_∞ has quadratic area growth, we can use the logarithmic cut-off trick exactly as in [30, page 54] to approximate the constant function 1 on Σ'_∞ by compactly supported functions in order to conclude that

$$\int_{\Sigma'_\infty} Q_{\Sigma'_\infty} d\mathcal{H}^2 \geq 0.$$

The strict dominant energy condition then implies that

$$\int_{\Sigma'_\infty} K_{\Sigma'_\infty} d\mathcal{H}^2 > 0, \quad (12)$$

where $K_{\Sigma'_\infty}$ denotes the Gauss curvature. On the other hand, just as in [27], the estimate $(g_\infty)_{ij}(x') - \delta_{ij} = O^2(|x'|^{-1})$ implies that the geodesic curvature of $\partial(\Sigma'_\infty \cap C_r)$ is $\kappa = \frac{1}{r} + O(r^{-2})$ while the length of $\partial(\Sigma'_\infty \cap C_r)$ is $2\pi r + O(1)$. The Gauss-Bonnet Theorem tells us that

$$\int_{\Sigma'_\infty \cap C_r} K_{\Sigma'_\infty} d\mathcal{H}^2 = 2\pi\chi(\Sigma'_\infty \cap C_r) - \int_{\partial(\Sigma'_\infty \cap C_r)} \kappa d\mathcal{H}^1.$$

Combined with (12) and the asymptotics of $\partial(\Sigma'_\infty \cap C_r)$ described above, for large r , we obtain

$$0 < 2\pi\chi(\Sigma'_\infty \cap C_r) - 2\pi.$$

Since $\Sigma'_\infty \setminus C_r$ is a graph for large r , we know that $\Sigma'_\infty \cap C_r$ is connected, yielding a contradiction. \square

5 The case $3 < n < 8$

Let $3 < n < 8$. We suppose that Theorem 1 holds in $n - 1$ dimensions and that it fails for an n -dimensional initial data set (M, g, k) as in Section 3. For the reasons described in the introduction, the argument here is substantially different from the proof in the time-symmetric case.

5.1 The functional \mathcal{F}

In this section we introduce a functional \mathcal{F} that will be essential for our proof. In order to motivate the definition of \mathcal{F} , consider the time-symmetric case when $k = 0$ so that the MOTS $\{\Sigma_{\rho,h}\}_{|h|\leq\Lambda}$ constructed in Section 3.2 are minimal hypersurfaces. An important step in the proof of the Riemannian positive mass theorem when $3 < n < 8$ [27] is to pick h_ρ such that Σ_{ρ,h_ρ} has least area in this family. Suppose for a moment that the family $\{\Sigma_{\rho,h}\}_{|h|\leq|\Lambda|}$ is actually a $C^{3,\alpha}$ foliation of minimal hypersurfaces with a first-order deformation vector field $X = \varphi\nu + \hat{X}$ that is equal

to ∂_n at ∂C_ρ . Then

$$\begin{aligned}
\frac{d}{dh} \mathcal{H}^{n-1}(\Sigma_{\rho,h}) &= \int_{\Sigma_{\rho,h}} (\operatorname{div}_{\Sigma_{\rho,h}} X) d\mathcal{H}^{n-1} \\
&= \int_{\Sigma_{\rho,h}} \operatorname{div}_{\Sigma_{\rho,h}} (\varphi \nu + \hat{X}) d\mathcal{H}^{n-1} \\
&= \int_{\Sigma_{\rho,h}} (H\varphi + \operatorname{div}_{\Sigma_{\rho,h}} \hat{X}) d\mathcal{H}^{n-1} \\
&= \int_{\Sigma_{\rho,h}} (\operatorname{div}_{\Sigma_{\rho,h}} \hat{X}) d\mathcal{H}^{n-1} \\
&= \int_{\partial\Sigma_{\rho,h}} \langle \hat{X}, \eta \rangle d\mathcal{H}^{n-2} \\
&= \int_{\partial\Sigma_{\rho,h}} \langle \partial_n, \eta \rangle d\mathcal{H}^{n-2}.
\end{aligned}$$

If $h_\rho \in (-\Lambda, \Lambda)$ minimizes areas as described above, then the *first derivative* in h of the integral above is nonnegative. This, along with the stability of Σ_{ρ,h_ρ} among deformations that keep the boundary fixed, is all that is needed to finish the proof in the time-symmetric case [27].

We now return to the general case. Instead of using the area functional, which is not adapted for application to MOTS, we will build our proof around the functional described below.

Definition 8. Let Σ be a compact hypersurface in M whose boundary lies on some coordinate cylinder ∂C_r . We let

$$\mathcal{F}(\Sigma) = \int_{\partial\Sigma} \langle \partial_n, \eta \rangle d\mathcal{H}^{n-2} \quad (13)$$

where η is the outward unit normal of $\partial\Sigma$ in Σ . Note that, using harmonic asymptotics, one can easily see that

$$\mathcal{F}(\Sigma) = \int_{\partial\Sigma} u^{\frac{2(n-1)}{n-2}} \eta_0^n d\mathcal{H}_0^{n-2} \quad (14)$$

where η_0^n is the n -th component of the unit normal η_0 computed using the Euclidean metric, and \mathcal{H}_0^{n-1} denotes Euclidean Hausdorff measure.

The barrier planes $\{x_n = \pm\Lambda\}$ give us a sign on $\mathcal{F}(\Sigma_{\rho,\pm\Lambda})$:

Lemma 11. *For any ρ sufficiently large,*

$$\mathcal{F}(\Sigma_{\rho,-\Lambda}) < 0 < \mathcal{F}(\Sigma_{\rho,\Lambda}).$$

Proof. From Lemma 6 we know that $\Sigma_{\rho,\Lambda}$ lies below the plane $\{x^n = \Lambda\}$ in C_ρ . The strong maximum principle Lemma 4 implies that they cannot meet tangentially at their common boundary $\Gamma_{\rho,\Lambda}$. Hence the Euclidean outward unit normal of $\partial\Sigma_{\rho,\Lambda}$ in $\Sigma_{\rho,\Lambda}$ satisfies $\eta_0^n > 0$. The inequality $\mathcal{F}(\Sigma_{\rho,\Lambda}) > 0$ then follows from (14). The proof that $\mathcal{F}(\Sigma_{\rho,-\Lambda}) < 0$ is analogous. \square

Recall the definition of jump heights from Section 3.3.

Lemma 12. *The function $h \mapsto \mathcal{F}(\Sigma_{\rho,h})$ is continuous at every $h_0 \in [-\Lambda, \Lambda]$ that is not a jump height. If $h_0 \in [-\Lambda, \Lambda]$ is a jump height, then*

$$\lim_{h \nearrow h_0} \mathcal{F}(\Sigma_{\rho,h}) \geq \mathcal{F}(\Sigma_{\rho,h_0}) \geq \lim_{h \searrow h_0} \mathcal{F}(\Sigma_{\rho,h}),$$

where both limits exist, and at least one of the inequalities above is strict. In other words, there must be a downward jump discontinuity at every jump height.

Proof. Let $h_0 \in [-\Lambda, \Lambda]$. Then by Lemma 7, $\lim_{h \nearrow h_0} \mathcal{F}(\Sigma_{\rho,h}) = \mathcal{F}(\underline{\Sigma}_{\rho,h_0})$ and $\lim_{h \searrow h_0} \mathcal{F}(\Sigma_{\rho,h}) = \mathcal{F}(\bar{\Sigma}_{\rho,h_0})$. By definition, if h_0 is not a jump height, then both of these limits must equal $\mathcal{F}(\Sigma_{\rho,h_0})$.

Let h_0 be a jump height. Since the family $\{\Sigma_{\rho,h}\}_{|h| \leq \Lambda}$ is ordered, it is clear that $\underline{\Sigma}_{\rho,h_0}$ lies beneath Σ_{ρ,h_0} , which lies beneath $\bar{\Sigma}_{\rho,h_0}$. Since they all share the common boundary Γ_{ρ,h_0} , we have $(\underline{\eta})_0^n \geq \eta_0^n \geq (\bar{\eta})_0^n$, where $\underline{\eta}, \eta, \bar{\eta}$ are the outward normals of Γ_{ρ,h_0} in $\underline{\Sigma}_{\rho,h_0}$, Σ_{ρ,h_0} , and $\bar{\Sigma}_{\rho,h_0}$, respectively. Since h_0 is jump height, $\underline{\Sigma}_{\rho,h_0} \neq \bar{\Sigma}_{\rho,h_0}$, so the strong maximum principle (Lemma 4) implies that at least one of the above inequalities is strict. By the definition of \mathcal{F} in (14),

$$\mathcal{F}(\underline{\Sigma}_{\rho,h_0}) \geq \mathcal{F}(\Sigma_{\rho,h_0}) \geq \mathcal{F}(\bar{\Sigma}_{\rho,h_0}),$$

where at least one of the inequalities is strict. \square

We now compute the first variation of \mathcal{F} . In view of Lemmas 11 and 12 we may hope to find $h_\rho \in (-\Lambda, \Lambda)$ such that the derivative of $h \rightarrow \mathcal{F}(\Sigma_{\rho,h})$ at h_ρ (defined in a suitably weak sense) is nonnegative.

Proposition 13. *Let Σ be a compact hypersurface with unit normal ν in M whose boundary lies on some ∂C_r . Let X be a C^1 vector field along Σ that is tangent to ∂C_r along $\partial\Sigma$. Let Z be a vector field of M such that $Z = \partial_n$ along ∂C_r . Then*

$$D\mathcal{F}|_\Sigma(X) = \int_{\partial\Sigma} \langle \phi \nabla \varphi + G(X), \eta \rangle d\mathcal{H}^{n-2} \quad (15)$$

where

$$G(X) = D_X Z - D_{\hat{Z}} \hat{X} + (\varphi H + \operatorname{div}_\Sigma \hat{X}) \hat{Z} - \phi S(\hat{X}) - \varphi S(\hat{Z}) \quad (16)$$

and where $X = \varphi\nu + \hat{X}$ and $Z = \phi\nu + \hat{Z}$ are the decompositions of X and Z into normal and tangential parts along Σ .

Proof. Note that $\mathcal{F}(\Sigma) = \int_{\partial\Sigma} \langle Z, \eta \rangle d\mathcal{H}^{n-2}$. Let e_1, \dots, e_{n-2} be a local orthonormal frame for the tangent space of $\partial\Sigma$. We can differentiate Z , the outward unit normal η of $\partial\Sigma$ in Σ , and the induced measure on $\partial\Sigma$ to obtain

$$D\mathcal{F}|_{\Sigma}(X) = \int_{\partial\Sigma} \left[\langle D_X Z, \eta \rangle + \left\langle Z, \langle D_{\eta} X, \nu \rangle \nu - \sum_{i=1}^{n-2} \langle D_{e_i} X, \eta \rangle e_i \right\rangle + \langle Z, \eta \rangle \operatorname{div}_{\partial\Sigma} X \right] d\mathcal{H}^{n-2}. \quad (17)$$

(The derivative of η is computed by differentiating the orthogonality relations.) Along $\partial\Sigma$, we introduce the decomposition $\hat{Z} = \psi\eta + Z^{\partial}$ into components that are normal and tangential to $\partial\Sigma$. The second term in the integrand of (17) is

$$\begin{aligned} \left\langle Z, \langle D_{\eta} X, \nu \rangle \nu - \sum_{i=1}^{n-2} \langle D_{e_i} X, \eta \rangle e_i \right\rangle &= \langle Z, \nu \rangle \langle D_{\eta} X, \nu \rangle - \sum_{i=1}^{n-2} \langle Z, e_i \rangle \langle D_{e_i} X, \eta \rangle \\ &= \phi \langle D_{\eta}(\varphi\nu + \hat{X}), \nu \rangle - \langle D_{Z^{\partial}} X, \eta \rangle \\ &= \phi(\nabla_{\eta}\varphi + \langle D_{\eta}\hat{X}, \nu \rangle) - \langle D_{Z^{\partial}} X, \eta \rangle \\ &= \langle \phi\nabla\varphi, \eta \rangle - \langle \phi S(\hat{X}), \eta \rangle - \langle D_{Z^{\partial}} X, \eta \rangle. \end{aligned} \quad (18)$$

The third term in the integrand of (17) is

$$\begin{aligned} \langle Z, \eta \rangle \operatorname{div}_{\partial\Sigma} X &= \langle Z, \eta \rangle (\operatorname{div}_{\Sigma} X - \langle D_{\eta} X, \eta \rangle) \\ &= \langle \hat{Z}, \eta \rangle (\varphi H + \operatorname{div}_{\Sigma} \hat{X}) - \psi \langle D_{\eta} X, \eta \rangle \\ &= \langle (\varphi H + \operatorname{div}_{\Sigma} \hat{X})\hat{Z}, \eta \rangle - \langle D_{\psi\eta} X, \eta \rangle. \end{aligned} \quad (19)$$

Notice that the first term in the integrand of (17), the first two terms of (18) and the first term of (19) combine to give

$$\langle \phi\nabla\varphi + D_X Z + (\varphi H + \operatorname{div}_{\Sigma} \hat{X})\hat{Z} - \phi S(\hat{X}), \eta \rangle. \quad (20)$$

The remaining two terms, which are the last term of (18) and the last term of (19), combine to give

$$\begin{aligned} -\langle D_{Z^{\partial}} X, \eta \rangle - \langle D_{\psi\eta} X, \eta \rangle &= -\langle D_{\hat{Z}} X, \eta \rangle \\ &= -\langle D_{\hat{Z}}(\varphi\nu + \hat{X}), \eta \rangle \\ &= -\langle \varphi D_{\hat{Z}}\nu + D_{\hat{Z}}\hat{X}, \eta \rangle \\ &= -\langle \varphi S(\hat{Z}) + D_{\hat{Z}}\hat{X}, \eta \rangle. \end{aligned} \quad (21)$$

The result follows from combining (20) and (21). \square

5.2 Height picking and stability

The following lemma, whose proof we defer to Section 5.4, will stand in for the geometric inequality (1) that was available in the time-symmetric case.

Lemma 14. *Let $\Sigma'_{\rho,h}$ denote the component of $\Sigma_{\rho,h}$ that contains the boundary $\Gamma_{\rho,h}$. For every large ρ there exists $h_\rho \in (-\Lambda, \Lambda)$ and a C^2 vector field X along Σ'_{ρ,h_ρ} that is equal to ∂_n along $\partial\Sigma'_{\rho,h_\rho} = \Gamma_{\rho,h_\rho}$ such that $\varphi = \langle X, \nu \rangle > 0$,*

$$D\theta^+|_{\Sigma'_{\rho,h_\rho}}(X) = 0, \quad (22)$$

and

$$D\mathcal{F}|_{\Sigma'_{\rho,h_\rho}}(X) \geq 0. \quad (23)$$

The proof of this lemma would be straightforward if the path $h \mapsto \Sigma_{\rho,h}$ of $C^{3,\alpha}$ hypersurfaces were differentiable in h (and if the $\Sigma_{\rho,h}$ were connected). By Lemma 11, we could find h_ρ such that

$$\frac{d}{dh}\mathcal{F}(\Sigma_{\rho,h})\Big|_{h=h_\rho} \geq 0.$$

We would then choose X to be the first-order deformation field of the family $\Sigma_{\rho,h}$ at $h = h_\rho$. The preceding inequality would turn into (23), and the fact that each $\Sigma_{\rho,h}$ is a MOTS would lead to (22). Unfortunately, $\Sigma_{\rho,h}$ need not be differentiable in h . In general, the family $\Sigma_{\rho,h}$ must contain jumps for topological reasons. From Lemma 12 we know that $\mathcal{F}(\Sigma_{\rho,h})$ can only jump down at a jump height, so the presence of jumps does not cause problems for finding h_ρ as described above. However, even in the absence of jumps, the lack of differentiability in h presents a technical challenge.

Notation. For the remainder of this section, we will abbreviate Σ'_{ρ,h_ρ} by Σ_ρ .

Lemma 14 allows us to conclude the following stability-like property, which the reader should compare to Proposition 3.

Lemma 15. *Let ρ be sufficiently large. Let X and φ be as in the statement of Lemma 14. For every C^1 function v on Σ_ρ that is equal to $\phi = \langle \partial_n, \nu \rangle$ along $\partial\Sigma_\rho$ we have that*

$$\int_{\Sigma_\rho} (|\nabla v|^2 + Qv^2) d\mathcal{H}^{n-1} + \int_{\partial\Sigma_\rho} \langle \bar{G}(X), \eta \rangle d\mathcal{H}^{n-2} \geq 0 \quad (24)$$

where

$$\bar{G}(X) = G(X) + \phi\varphi W. \quad (25)$$

Proof. We begin by following the argument in [21]. Using equation (3) and the positivity of φ , we compute that

$$\begin{aligned} D\theta^+|_{\Sigma_\rho}(X) &= -\Delta\varphi + 2\langle W, \nabla\varphi \rangle + (\operatorname{div} W - |W|^2 + Q)\varphi \\ &= -\Delta\varphi + |\nabla \log \varphi|^2\varphi - |W - \nabla \log \varphi|^2\varphi + (\operatorname{div} W + Q)\varphi \\ &= -(\Delta \log \varphi)\varphi - |W - \nabla \log \varphi|^2\varphi + (\operatorname{div} W + Q)\varphi \\ &= [\operatorname{div}(W - \nabla \log \varphi)]\varphi - |W - \nabla \log \varphi|^2\varphi + Q\varphi. \end{aligned}$$

Let $v \in C^1(\Sigma)$ be equal to ϕ along $\partial\Sigma_\rho$. We multiply the above equation by $v^2\varphi^{-1}$ to obtain

$$\begin{aligned} v^2\varphi^{-1}D\theta^+|_{\Sigma_\rho}(X) &= [\operatorname{div}(W - \nabla \log \varphi)]v^2 - |W - \nabla \log \varphi|^2v^2 + Qv^2 \\ &= \operatorname{div}(v^2(W - \nabla \log \varphi)) - \langle W - \nabla \log \varphi, 2v\nabla v \rangle \\ &\quad - |W - \nabla \log \varphi|^2v^2 + Qv^2 \\ &= \operatorname{div}(v^2(W - \nabla \log \varphi)) \\ &\quad + |(W - \nabla \log \varphi)v|^2 + |\nabla v|^2 - |(W - \nabla \log \varphi)v + \nabla v|^2 \\ &\quad - |W - \nabla \log \varphi|^2v^2 + Qv^2 \\ &= \operatorname{div}(v^2(W - \nabla \log \varphi)) + |\nabla v|^2 + Qv^2 \\ &\quad - |(W - \nabla \log \varphi)v + \nabla v|^2. \end{aligned} \tag{26}$$

Together with equation (22), this implies that

$$0 \leq |\nabla v|^2 + Qv^2 + \operatorname{div}(v^2(W - \nabla \log \varphi)).$$

Using that $v = \phi = \varphi$ along $\partial\Sigma_\rho$, we estimate

$$\begin{aligned} 0 &\leq \int_{\Sigma_\rho} (|\nabla v|^2 + Qv^2) d\mathcal{H}^{n-1} + \int_{\partial\Sigma_\rho} \langle v^2(W - \nabla \log \varphi), \eta \rangle d\mathcal{H}^{n-2} \\ &= \int_{\Sigma_\rho} (|\nabla v|^2 + Qv^2) d\mathcal{H}^{n-1} + \int_{\partial\Sigma_\rho} \langle \phi\varphi W - \phi\nabla\varphi, \eta \rangle d\mathcal{H}^{n-2} \\ &= \int_{\Sigma_\rho} (|\nabla v|^2 + Qv^2) d\mathcal{H}^{n-1} + \int_{\partial\Sigma_\rho} \langle \bar{G}(X), \eta \rangle d\mathcal{H}^{n-2} - D\mathcal{F}|_{\Sigma_\rho}(X) \\ &\leq \int_{\Sigma_\rho} (|\nabla v|^2 + Qv^2) d\mathcal{H}^{n-1} + \int_{\partial\Sigma_\rho} \langle \bar{G}(X), \eta \rangle d\mathcal{H}^{n-2} \end{aligned}$$

where we used (15) and (23) in the third and fourth lines, respectively. \square

5.3 Analysis of the complete MOTS limit

For the remainder of the proof of Theorem 1 we follow the strategy for the time-symmetric case treated in [27] closely. By Lemma 8, there is a sequence $\rho_j \rightarrow \infty$

such that the Σ_{ρ_j} converge locally in $C^{3,\alpha}$ to a complete MOTS Σ_∞ that has the properties described in Lemma 8.

Lemma 10 is strong enough to show that Σ_∞ is conformal to a scalar-flat asymptotically flat manifold but it does not provide sufficient control on the change of mass effected by this conformal change. For that, we need to allow test functions that are asymptotic to 1. To justify the use of such test functions we need some uniform control on the Σ_ρ 's as in the following lemma.

Lemma 16. *Let Z be a C^2 vector field on M that is equal to ∂_n outside a compact set and let $Z = \phi\nu + \hat{Z}$ be the decomposition of Z into normal and tangential part along Σ_ρ . The following estimate holds uniformly in ρ large.*

$$\left(\int_{\Sigma_\rho \setminus C_r} (|\nabla \phi|^2 + Q_{\Sigma_\rho} \phi^2) d\mathcal{H}^{n-1} + \int_{\partial(\Sigma_\rho \setminus C_r)} \langle \bar{G}(Z), \eta \rangle d\mathcal{H}^{n-2} \right) = O(r^{-1}). \quad (27)$$

Proof. The estimates

$$D\theta^+|_{\Sigma_\rho}(Z) = O(|x|^{-n}) \quad (28)$$

$$D\mathcal{F}|_{\Sigma_\rho \cap C_r}(Z) = O(r^{-1}) \quad (29)$$

hold uniformly in ρ large. To see this, we use the harmonic asymptotics and formula (7) to obtain that

$$H = u^{\frac{-2}{n-2}} \left(H_0 + \frac{2(n-1)}{n-2} u^{-1} \nabla_{\nu_0} u \right)$$

where H_0 and ν_0 are the mean curvature and upward unit normal with respect to the Euclidean metric. Clearly, H_0 and ν_0 do not change under vertical translation, and H_0 is bounded. Therefore, the decay of u implies that $DH|_{\Sigma_\rho}(Z) = O(|x|^{-n})$. The $\text{tr}_{\Sigma_\rho} k$ term is easier to handle, giving us (28). To derive (29), we simply use (14), the fact that ν_0 does not change under vertical translation, and the decay of u .

We vary Σ_ρ in the direction Z , which is just vertical translation outside a compact set. We repeat the argument from the beginning of the proof of Lemma 15, except that we use $\Sigma_\rho \setminus C_r$ instead of Σ_ρ , the vector field Z instead of X , the function ϕ instead of φ , and that we take $v = \phi$. On $\Sigma_\rho \setminus C_r$, equation (26) becomes

$$\phi D\theta^+|_{\Sigma_\rho}(Z) = \text{div}(\phi^2 W - \phi \nabla \phi) + |\nabla \phi|^2 + Q\phi^2 - |W|^2 \phi^2.$$

Therefore,

$$\begin{aligned}
& \int_{\Sigma_\rho \setminus C_r} (|\nabla \phi|^2 + Q\phi^2) d\mathcal{H}^{n-1} + \int_{\partial(\Sigma_\rho \setminus C_r)} \langle \bar{G}(Z), \eta \rangle d\mathcal{H}^{n-2} \\
&= \int_{\Sigma_\rho \setminus C_r} (|\nabla \phi|^2 + Q\phi^2) d\mathcal{H}^{n-1} + \int_{\partial(\Sigma_\rho \setminus C_r)} \langle \phi^2 W - \phi \nabla \phi, \eta \rangle d\mathcal{H}^{n-2} \\
&\quad + D\mathcal{F}|_{\Sigma_\rho}(Z) - D\mathcal{F}|_{\Sigma_\rho \cap C_r}(Z) \\
&= \int_{\Sigma_\rho \setminus C_r} (\phi D\theta^+|_{\Sigma_\rho}(Z) + |W|^2 \phi^2) d\mathcal{H}^{n-1} \\
&= O(r^{-1})
\end{aligned}$$

where the last line follows from the (28) and (29), decay of W , and volume control coming from the almost minimizing property of Σ_ρ . \square

The following lemma is the analytic consequence of our careful height-picking in Section 5.2.

Lemma 17. *Let v be a function on Σ_∞ such that $v - 1 \in W_{\frac{3-n}{2}}^{1,2}(\Sigma_\infty)$. Then*

$$\int_{\Sigma_\infty} (|\nabla v|^2 + Q_{\Sigma_\infty} v^2) d\mathcal{H}^{n-1} \geq 0. \quad (30)$$

Proof. Following the notation of Lemma 8, we use coordinates x' on $\Sigma_\infty \setminus B$ where B is a large compact subset of M . Lemma 8 gives that

$$Q_{\Sigma_\infty} = \frac{1}{2}R_{\Sigma_\infty} - \mu - J(\nu_{\Sigma_\infty}) - \frac{1}{2}|k_{\Sigma_\infty} + B_{\Sigma_\infty}|^2 = O(|x'|^{-n}).$$

Using that the volume of Σ_∞ grows like that of \mathbb{R}^{n-1} , it follows that

$$\int_{\Sigma_\infty} (|\nabla v|^2 + |Q_{\Sigma_\infty}|v^2) d\mathcal{H}^{n-1} < \infty \quad (31)$$

provided that $v - 1 \in W_{\frac{3-n}{2}}^{1,2}(\Sigma_\infty)$.

Below we abbreviate $\Sigma_{\rho_j} = \Sigma'_{\rho_j, h_{\rho_j}}$ by Σ_j . We also use a subscript j on geometric quantities to indicate that they are computed with respect to Σ_j . We use η_j to denote the outward unit normal of $\partial(\Sigma_j \cap C_r)$ in Σ_j for large $r < \rho_j$ such that ∂C_r is transverse to each Σ_j . Fix a C^2 vector field Z on M that agrees with ∂_n outside a compact set. Let $\phi_\infty = \langle \nu_{\Sigma_\infty}, Z \rangle$ and $\phi_j = \langle \nu_{\Sigma_j}, Z \rangle$. It follows from Lemma 8 that $\phi_\infty - 1 = O^1(|x'|^{2-n}) \in W_{(3-n)/2}^{1,2}(\Sigma_\infty)$ and that $\bar{G}_\infty(Z) = O(|x'|^{1-n})$ where \bar{G}_∞ is as in (25). We have that

$$\begin{aligned}
& \int_{\Sigma_\infty} (|\nabla \phi_\infty|^2 + Q_\infty \phi_\infty^2) d\mathcal{H}^{n-1} \\
&= \lim_{r \rightarrow \infty} \int_{\Sigma_\infty \cap C_r} (|\nabla \phi_\infty|^2 + Q_\infty \phi_\infty^2) d\mathcal{H}^{n-1} \\
&= \lim_{r \rightarrow \infty} \left(\int_{\Sigma_\infty \cap C_r} (|\nabla \phi_\infty|^2 + Q_\infty \phi_\infty^2) d\mathcal{H}^{n-1} + \int_{\partial(\Sigma_\infty \cap C_r)} \langle \bar{G}_\infty(Z), \eta_\infty \rangle d\mathcal{H}^{n-2} \right) \\
&= \lim_{r \rightarrow \infty} \lim_{j \rightarrow \infty} \left(\int_{\Sigma_j \cap C_r} (|\nabla \phi_j|^2 + Q_j \phi_j^2) d\mathcal{H}^{n-1} + \int_{\partial(\Sigma_j \cap C_r)} \langle \bar{G}_j(Z), \eta_j \rangle d\mathcal{H}^{n-2} \right) \\
&= \lim_{j \rightarrow \infty} \left(\int_{\Sigma_j} (|\nabla \phi_j|^2 + Q_j \phi_j^2) d\mathcal{H}^{n-1} + \int_{\partial \Sigma_j} \langle \bar{G}_j(Z), \eta_j \rangle d\mathcal{H}^{n-2} \right) \\
&\geq 0,
\end{aligned}$$

where the last inequality follows from Lemma 15, and the fourth equality follows from Lemma 16. It follows that (30) holds for $v = \phi_\infty$. Moreover, since the argument works for any Z that equals ∂_n outside a compact set, (30) holds for all test functions that agree with ϕ_∞ outside a compact set.

We now argue by density that (30) holds for any function v such that $v - \phi_\infty \in W_{(3-n)/2}^{1,2}(\Sigma_\infty)$. Note that $C_c^{3,\alpha}(\Sigma_\infty)$ is dense in $W_{(3-n)/2}^{1,2}(\Sigma_\infty)$. Let $v_i - \phi_\infty$ be a sequence of functions in $C_c^{3,\alpha}(\Sigma_\infty)$ that converges to $v - \phi_\infty$ in $W_{(3-n)/2}^{1,2}(\Sigma_\infty)$. It is straightforward to check that

$$\begin{aligned}
0 &\leq \liminf_{i \rightarrow \infty} \int_{\Sigma_\infty} (|\nabla v_i|^2 + Q_\infty v_i^2) d\mathcal{H}^{n-1} \\
&= \liminf_{i \rightarrow \infty} \int_{\Sigma_\infty} [(|\nabla v|^2 + Q_\infty v^2) + (2\nabla v \cdot \nabla(v_i - v) + 2Q_\infty v(v_i - v)) \\
&\quad + (|\nabla(v_i - v)|^2 + Q_\infty(v_i - v)^2)] d\mathcal{H}^{n-1} \\
&= \int_{\Sigma_\infty} (|\nabla v|^2 + Q_\infty v^2) d\mathcal{H}^{n-1}
\end{aligned}$$

where the cross terms vanish because of (31). This implies the result since $\phi_\infty - 1 \in W_{(3-n)/2}^{1,2}(\Sigma_\infty)$. \square

We are now ready to derive a contradiction to our induction hypothesis. Without loss of generality, we throw away all compact components of Σ_∞ , noting that Lemmas 10 and 17 still hold on the remaining asymptotically flat component. The strict dominant energy condition implies that $Q_\infty < \frac{1}{2}R_{\Sigma_\infty}$. Lemma 10 implies that for any nonzero $v \in W_{(3-n)/2}^{1,2}(\Sigma_\infty)$, we have

$$\int_{\Sigma_\infty} (|\nabla v|^2 + \frac{1}{2}R_{\Sigma_\infty} v^2) d\mathcal{H}^{n-1} > 0.$$

Using that $n > 3$, we have

$$\int_{\Sigma_\infty} \left(|\nabla v|^2 + \frac{n-3}{4(n-2)} R_{\Sigma_\infty} v^2 \right) d\mathcal{H}^{n-1} > 0 \quad (32)$$

for all nonzero $v \in W_{(3-n)/2}^{1,2}(\Sigma_\infty)$. This implies that the conformal Laplacian

$$\Delta_{\Sigma_\infty} - \frac{n-3}{4(n-2)} R_{\Sigma_\infty} : W_{\frac{3-n}{2}}^{1,2}(\Sigma_\infty) \rightarrow W_{\frac{-1-n}{2}}^{-1,2}(\Sigma_\infty)$$

is an isomorphism. In particular, we can find some nonzero $v \in W_{(3-n)/2}^{1,2}(\Sigma_\infty)$ such that

$$\Delta_{\Sigma_\infty} v - \frac{n-3}{4(n-2)} R_{\Sigma_\infty} v = \frac{n-3}{4(n-2)} R_{\Sigma_\infty}.$$

Setting $w = 1 + v$, we have that

$$\Delta_{\Sigma_\infty} w - \frac{n-3}{4(n-2)} R_{\Sigma_\infty} w = 0.$$

Note that $w \in C_{loc}^{2,\alpha}$ by elliptic regularity. By [24, Theorem 2], we have that $w(x') = 1 + O^{2,\alpha}(|x'|^{3-n})$. Let g_∞ be the induced metric on Σ_∞ . Applying (6) in dimension $n-1$, the conformal metric $w^{\frac{4}{n-3}} g_\infty$ on Σ_∞ is asymptotically flat with zero scalar curvature. Using (8) in dimension $n-1$, this metric has energy

$$\begin{aligned} E(w^{\frac{4}{n-3}} g_\infty) &= E(g_\infty) - \frac{2}{(n-3)\omega_{n-2}} \lim_{r \rightarrow \infty} \int_{\partial(\Sigma_\infty \cap C_r)} w \nabla_\nu w d\mathcal{H}^{n-1} \\ &= 0 - \frac{2}{(n-3)\omega_{n-2}} \int_{\Sigma_\infty} (|\nabla w|^2 + w \Delta w) d\mathcal{H}^{n-1} \\ &= \frac{-2}{(n-3)\omega_{n-2}} \int_{\Sigma_\infty} \left(|\nabla w|^2 + \frac{n-3}{4(n-2)} R_{\Sigma_\infty} w^2 \right) d\mathcal{H}^{n-1}. \end{aligned}$$

Since $w - 1 \in W_{(3-n)/2}^{1,2}(\Sigma_\infty)$ is nonzero, we can apply Lemma 17 and the short argument used to derive (32) to conclude that

$$\int_{\Sigma_\infty} \left(|\nabla w|^2 + \frac{n-3}{4(n-2)} R_{\Sigma_\infty} w^2 \right) d\mathcal{H}^{n-1} > 0.$$

It follows that $E(w^{\frac{4}{n-3}} g_\infty) < 0$. This contradicts the time-symmetric case of Theorem 1 in dimension $n-1$. In particular, it contradicts our induction hypothesis.

5.4 Proof of Lemma 14

We define

$$h_\rho = \inf\{h \in [-\Lambda, \Lambda] \mid \mathcal{F}(\Sigma_{\rho,h}) > 0\}. \quad (33)$$

Our goal for this section is to prove that this choice of h_ρ satisfies the conclusion of Lemma 14. For now let us assume that Σ_{ρ,h_ρ} is connected.

Using Lemmas 11 and 12 it is easy to see that h_ρ exists, lies in $(-\Lambda, \Lambda)$, is not a jump height, and that $\mathcal{F}(\Sigma_{\rho, h_\rho}) = 0$. Although the path $h \mapsto \Sigma_{\rho, h}$ of $C^{3,\alpha}$ hypersurfaces need not be differentiable in h at h_ρ , the continuity of the map $h \mapsto \Sigma_{\rho, h}$ at $h = h_\rho$ allows us to use the inverse function theorem to describe the family $\Sigma_{\rho, h}$ precisely for h near h_ρ , as we will see below. We note that the work of B. White [36] on the moduli space of minimal submanifolds with boundary in \mathbb{R}^n is useful for understanding the family $\Sigma_{\rho, h}$. However, since we do not need the full power of [36], we choose to use a simpler approach similar to that of [19] in our Case 2 below.

Since Σ_{ρ, h_ρ} is transverse to ∂C_ρ , we can find a $C^{2,\alpha}$ vector field ∂_τ such that $\langle \partial_\tau, \nu \rangle > 0$ along Σ_{ρ, h_ρ} , and $\partial_\tau = Z = \partial_n$ at Γ_{ρ, h_ρ} . By integrating ∂_τ , there exists a relatively open neighborhood U of Σ_{ρ, h_ρ} in C_ρ and a $C^{2,\alpha}$ diffeomorphism $F : \Sigma_{\rho, h_\rho} \times (-\delta, \delta) \rightarrow U$ such that $F(\cdot, 0)$ is the identity map and such that $F_*(\frac{\partial}{\partial \tau}) = \partial_\tau$. In particular, $F(\cdot, \tau)$ maps $\partial \Sigma_{\rho, h_\rho} = \Gamma_{\rho, h_\rho}$ to $\Gamma(h_\rho + \tau)$ by vertical translation.

Every $C^{2,\alpha}$ function $w : \Sigma_{\rho, h_\rho} \rightarrow (-\delta, \delta)$ gives rise to a graph in $\Sigma_{\rho, h_\rho} \times (-\delta, \delta)$ whose image under the diffeomorphism F is a $C^{2,\alpha}$ hypersurface in U denoted by $\text{graph}[w]$ whose boundary lies on ∂C_ρ . Since h_ρ is not a jump height, Lemma 7 tells us that for each h sufficiently close to h_ρ the MOTS $\Sigma_{\rho, h}$ coincides with $\text{graph}[w]$ for a unique $C^{2,\alpha}$ function w . Moreover, the function w converges to 0 in $C^{2,\alpha}$ as h approaches h_ρ .

The operator $L_{\Sigma_{\rho, h_\rho}}$ on Σ_{ρ, h_ρ} defined by (4) has a principal Dirichlet eigenvalue, which is nonnegative because Σ_{ρ, h_ρ} is stable. We consider two cases.

Case 1: The principal eigenvalue of $L_{\Sigma_{\rho, h_\rho}}$ is positive.

Define the map³ $\Psi : C_0^{2,\alpha}(\Sigma_{\rho, h_\rho}) \times \mathbb{R} \rightarrow C^{0,\alpha}(\Sigma_{\rho, h_\rho}) \times \mathbb{R}$ by

$$\Psi(w, s) = (\theta_{\text{graph}[w+s]}^+, s).$$

Using (3) and (4), we see that

$$D\Psi|_{(0,0)}(w', s') = (L_{\Sigma_{\rho, h_\rho}} \langle (w' + s') \partial_\tau, \nu \rangle, s').$$

By assumption, $L_{\Sigma_{\rho, h_\rho}} : C_0^{2,\alpha} \rightarrow C^{0,\alpha}$ is an isomorphism. It follows that $D\Psi|_{(0,0)} : C_0^{2,\alpha}(\Sigma_{\rho, h_\rho}) \times \mathbb{R} \rightarrow C^{0,\alpha}(\Sigma_{\rho, h_\rho}) \times \mathbb{R}$ is an isomorphism as well. The inverse function theorem gives a C^1 function $\Phi : (-\epsilon, \epsilon) \rightarrow C_0^{2,\alpha}(\Sigma_{\rho, h_\rho})$ such that $\Phi(s)$ is the unique small solution of the equation $\Psi(\Phi(s), s) = (0, s)$. Its graph is the unique $C^{2,\alpha}$ nearby MOTS with boundary $\Gamma_{\rho, h_\rho+s}$. By continuity of $\Sigma_{\rho, h}$ at $h = h_\rho$ and uniqueness, we have that $\Sigma_{\rho, h_\rho+s} = \text{graph}[\Phi(s)+s]$ for small s . It follows that $h \mapsto \Sigma_{\rho, h}$ is a C^1 path

³Technically, it is only defined on some neighborhood of the origin.

into the space of $C^{2,\alpha}$ hypersurfaces near $h = h_\rho$ to which the straightforward argument described in Section 5.2 applies. Specifically, consider the $C^{2,\alpha}$ first-order deformation field $X = (\frac{d\Phi}{ds} + 1)\partial_\tau$ of the family $\Sigma_{\rho,h}$. Then $D\theta^+|_{\Sigma_{\rho,h_\rho}}(X) = \frac{d}{dh}\theta_{\Sigma_{\rho,h}}^+|_{h=h_\rho} = 0$ because each $\Sigma_{\rho,h}$ is a MOTS. That $D\mathcal{F}|_{\Sigma_{\rho,h_\rho}}(X) = \frac{d}{dh}\mathcal{F}(\Sigma_{\rho,h})|_{h=h_\rho} \geq 0$ follows from the construction of h_ρ . The normal component φ of X along $\Sigma_{\rho,h}$ is nonnegative and positive on $\Gamma_{\rho,h}$ by construction, and thus positive everywhere by the strong maximum principle applied to the linear equation $L_{\Sigma_{\rho,h_\rho}}\varphi = 0$. This completes the proof of Lemma 14 for Case 1.

Case 2: The principal eigenvalue of $L_{\Sigma_{\rho,h_\rho}}$ is zero.

In this case, since Σ_{ρ,h_ρ} is connected, $L_{\Sigma_{\rho,h_\rho}}$ has a one-dimensional kernel in $C_0^{2,\alpha}$ that is generated by a function that is positive away from the boundary. The same is then true for the adjoint $L_{\Sigma_{\rho,h_\rho}}^*$. We define the map $\Psi : C_0^{2,\alpha}(\Sigma_{\rho,h_\rho}) \times \mathbb{R}^2 \rightarrow C^{0,\alpha}(\Sigma_{\rho,h_\rho}) \times \mathbb{R}^2$ by

$$\Psi(w, \kappa, s) = \left(-\kappa + \theta_{\text{graph}[w+s]}^+, V(\text{graph}[w+s]), s \right)$$

where $V(\Sigma)$ denotes the signed volume of the region of C_ρ lying above Σ_{ρ,h_ρ} and below Σ . Then

$$D\Psi|_{(0,0,0)}(w', \kappa', s') = \left(L_{\Sigma_{\rho,h_\rho}} \langle (w' + s')\partial_\tau, \nu \rangle - \kappa', \int_{\Sigma_{\rho,h_\rho}} \langle (w' + s')\partial_\tau, \nu \rangle d\mathcal{H}^{n-1}, s' \right).$$

We claim that $D\Psi|_{(0,0,0)}$ is injective. If $D\Psi|_{(0,0,0)}(w', \kappa', s') = (0, 0, 0)$, then obviously $s' = 0$, and since the image of $C_0^{2,\alpha}$ under $L_{\Sigma_{\rho,h_\rho}}$ is orthogonal to the kernel of $L_{\Sigma_{\rho,h_\rho}}^*$, $\kappa' = 0$. Finally, $\langle w'\partial_\tau, \nu \rangle$ is in the kernel of $L_{\Sigma_{\rho,h_\rho}}$ and has zero integral, and is therefore zero, proving the claim. Since $D\Psi|_{(0,0,0)}$ has index zero, it is also an isomorphism. By the inverse function theorem, there exists a C^1 function

$$(\Phi_1, \Phi_2) : (\epsilon, \epsilon) \times (\epsilon, \epsilon) \rightarrow C_0^{2,\alpha}(\Sigma_{\rho,h_\rho}) \times \mathbb{R}$$

such that $(\Phi_1(\xi, s), \Phi_2(\xi, s))$ is the unique small solution of the equation

$$\Psi(\Phi_1(\xi, s), \Phi_2(\xi, s), s) = (0, \xi, s).$$

In particular, $\text{graph}[\Phi_1(\xi, s) + s]$ is the unique $C^{2,\alpha}$ -nearby constant θ^+ hypersurface whose boundary is $\Gamma_{\rho,h_\rho+s}$ and whose signed volume is ξ . Define $\xi(s) := V(\Sigma_{\rho,h_\rho+s})$. By continuity of $\Sigma_{\rho,h}$ at $h = h_\rho$ and the uniqueness, it must be the case that $\Sigma_{\rho,h_\rho+s} = \text{graph}[\Phi_1(\xi(s), s) + s]$ for small s . Note that since $\Sigma_{\rho,h_\rho+s}$ lies strictly above Σ_{ρ,h_ρ} , the function $\Phi_1(\xi(s), s) + s$ must be positive. The complication here is that we do not know that $\xi(s)$ depends on s in any nice way. Below, we will see that this does not matter.

By the construction of h_ρ , we can choose a sequence $s_k \searrow 0$ such that

$$\mathcal{F}(\Sigma_{\rho, h_\rho + s_k}) > 0.$$

Let us pass to a subsequence such that the unit vector in the direction of $(\xi(s_k), s_k)$ converges. We consider two subcases.

Case 2a: The ratio $\frac{\xi(s_k)}{s_k}$ converges to a finite number as $k \rightarrow \infty$.

The basic idea here is while we cannot take derivatives as in Case 1, we can take subsequential limits of difference quotients instead. The hypothesis of this subcase implies that $\frac{1}{s_k}[\Phi_1(\xi(s_k), s_k) + s_k]$ converges to a $C^{2,\alpha}$ function $\bar{\varphi}$. Therefore we can take the limit of $\frac{1}{s_k}\theta^+_{\text{graph}[\Phi_1(\xi(s_k), s_k) + s_k]} = 0$ as $k \rightarrow \infty$ to obtain

$$L_{\Sigma_{\rho, h_\rho}}(\langle \bar{\varphi} \partial_\tau, \nu \rangle) = 0.$$

We claim that the conclusion of Lemma 14 holds for $X = \bar{\varphi} \partial_\tau \in C^{2,\alpha}$. The previous equation tells us that $D\theta^+|_{\Sigma_{\rho, h_\rho}}(X) = 0$. Moreover, the normal component $\varphi = \bar{\varphi} \langle \partial_\tau, \nu \rangle$ is nonnegative and equal to $\langle Z, \nu \rangle > 0$ at Γ_{ρ, h_ρ} . In fact, we see that φ is positive by the strong maximum principle applied to the operator $L_{\Sigma_{\rho, h_\rho}}$. Finally, since $\mathcal{F}(\Sigma_{\rho, h_\rho + s_k}) > 0$, we have that

$$\begin{aligned} D\mathcal{F}|_{\Sigma_{\rho, h_\rho}}(X) &= \lim_{k \rightarrow \infty} \frac{1}{s_k} [\mathcal{F}(\text{graph}[\Phi_1(\xi(s_k), s_k) + s_k]) - \mathcal{F}(\Sigma_{\rho, h_\rho})] \\ &= \lim_{k \rightarrow \infty} \frac{1}{s_k} [\mathcal{F}(\Sigma_{\rho, h_\rho + s_k}) - \mathcal{F}(\Sigma_{\rho, h_\rho})] \\ &\geq 0. \end{aligned}$$

Case 2b: The ratio $\frac{s_k}{\xi(s_k)}$ converges to zero as $k \rightarrow \infty$.

We will show that this is impossible. The assumption implies that the quotients $\frac{1}{\xi(s_k)}[\Phi_1(\xi(s_k), s_k) + s_k]$ converge to a $C^{2,\alpha}$ function $\bar{\varphi}$ that vanishes along Γ_{ρ, h_ρ} . As in Case 2a, we conclude that

$$L_{\Sigma_{\rho, h_\rho}}(\langle \bar{\varphi} \partial_\tau, \nu \rangle) = 0.$$

Let $X = \bar{\varphi} \partial_\tau \in C^{2,\alpha}$. Its normal component $\varphi = \bar{\varphi} \langle \partial_\tau, \nu \rangle$ is nonnegative with zero boundary values. It follows from the strong maximum principle that either φ is identically zero, or else $\nabla_\eta \varphi < 0$ along Γ_{ρ, h_ρ} . Assume the latter. Taking the limit of

the equation $V(\Sigma_{\rho,h_\rho+s_k})/\xi(s_k) = 1$ we obtain $\int_{\Sigma_{\rho,h_\rho}} \varphi d\mathcal{H}^{n-1} = 1$. As in Case 2a,

$$\begin{aligned} D\mathcal{F}|_{\Sigma_{\rho,h_\rho}}(X) &= \lim_{k \rightarrow \infty} \frac{1}{\xi(s_k)} [\mathcal{F}(\text{graph}[\Phi_1(\xi(s_k), s_k) + s_k]) - \mathcal{F}(\Sigma_{\rho,h_\rho})] \\ &= \lim_{k \rightarrow \infty} \frac{1}{\xi(s_k)} [\mathcal{F}(\Sigma_{\rho,h_\rho+s_k}) - \mathcal{F}(\Sigma_{\rho,h_\rho})] \\ &\geq 0. \end{aligned}$$

On the other hand, only the second term in (17) contributes to $D\mathcal{F}|_{\Sigma_{\rho,h_\rho}}$ because X vanishes along Γ_{ρ,h_ρ} . Thus

$$\begin{aligned} D\mathcal{F}|_{\Sigma_{\rho,h_\rho}}(X) &= \int_{\Gamma_{\rho,h_\rho}} \langle Z, \langle D_\eta X, \nu \rangle \nu \rangle d\mathcal{H}^{n-2} \\ &= \int_{\Gamma_{\rho,h_\rho}} \phi \nabla_\eta \varphi d\mathcal{H}^{n-2} \\ &< 0. \end{aligned}$$

This contradiction shows that Case 2b cannot occur.

The proof of Lemma 14 in the case where Σ_{ρ,h_ρ} is connected is now finished. Assume now that Σ_{ρ,h_ρ} is not connected. Let $\Sigma'_{\rho,h}$ denote the component of $\Sigma_{\rho,h}$ that contains the boundary. Observe that $\Sigma'_{\rho,h}$ converges to Σ'_{ρ,h_ρ} in $C^{3,\alpha}$ as $h \rightarrow h_\rho$ because h_ρ is not a jump height. Moreover, $\mathcal{F}(\Sigma'_{\rho,h}) = \mathcal{F}(\Sigma_{\rho,h})$ for all $h \in [-\Lambda, \Lambda]$. From this it is easy to see that the above argument can be carried out with Σ'_{ρ,h_ρ} in place of $\Sigma_{\rho,h}$.

6 The density theorem

Recall that $\pi = k - (\text{tr}_g k)g$ and that

$$n \geq 3, \quad p > n, \quad q \in ((n-2)/2, n-2), \quad q_0 > 0 \quad \text{and} \quad \alpha \in (0, 1 - n/p].$$

It will be convenient to express initial data in terms of π rather than k . Abusing terminology slightly, we will refer to (M, g, π) as an initial data set in this section. We denote by g_E a fixed smooth symmetric $(0, 2)$ -tensor that coincides with the Euclidean metric on $M \setminus K \cong \mathbb{R}^n \setminus B$ throughout this section.

Theorem 18 (Density theorem). *Let (M, g, π) be an n -dimensional asymptotically flat initial data set of type (p, q, q_0, α) such that the dominant energy condition $\mu \geq |J|_g$ holds. Let $\epsilon > 0$. There are asymptotically flat initial data $(\bar{g}, \bar{\pi})$ with harmonic asymptotics and of type (p, q, q_0, α) on M such that*

$$\|g - \bar{g}\|_{W_{-q}^{2,p}} < \epsilon, \quad \|\pi - \bar{\pi}\|_{W_{-1-q}^{1,p}} < \epsilon, \quad |E - \bar{E}| < \epsilon, \quad |P - \bar{P}| < \epsilon,$$

and such that the strict dominant energy condition

$$\bar{\mu} > |\bar{J}|_{\bar{g}}$$

holds.

Remark. If we assume appropriate higher regularity for (M, g, π) , then $(M, \bar{g}, \bar{\pi})$ will have the same regularity. This follows from applying Schauder estimates throughout the proof. Our argument shows that $(\bar{g}, \bar{\pi})$ can be taken to be of type (p, q, q'_0, α) for any given $q'_0 > q_0$.

The proof of Theorem 18 consists of two general constructions. In Section 6.1 we deform (g, π) to initial data $(\hat{g}, \hat{\pi})$ such that $\hat{\mu} > (1 + \gamma)|\hat{J}|_{\hat{g}}$ for some $\gamma > 0$. In Section 6.2 we apply a cut-off argument to perturb $(\hat{g}, \hat{\pi})$ to harmonic asymptotics as in [11, Theorem 1] and argue that we can preserve the strict dominant energy condition in the process.

The ADM energy and linear momentum are continuous on the space of asymptotically flat initial data sets in the following sense.

Proposition 19 (*Cf.* [22, Proposition 2.4]). *Let (g, π) and $(\bar{g}, \bar{\pi})$ be asymptotically flat initial data of type (p, q, q_0, α) . Let $\epsilon > 0$. There exists $\delta > 0$ depending only on ϵ, n, p, q, q_0 , $\|(g - g_{\mathbb{E}}, \pi)\|_{W_{-q}^{2,p} \times W_{-q-1}^{1,p}}$, and $\|(\mu, J) - (\bar{\mu}, \bar{J})\|_{L_{-n-q_0/2}^1}$ such that if*

$$\|g - \bar{g}\|_{W_{-q}^{2,p}} \leq \delta \quad \text{and} \quad \|\pi - \bar{\pi}\|_{W_{-1-q}^{1,p}} \leq \delta,$$

then

$$|E - \bar{E}| < \epsilon \quad \text{and} \quad |P - \bar{P}| < \epsilon.$$

The proof of this fact is now standard and goes back to [31, p. 50] (for E only) and [11, p. 198] in the case of vacuum data. We include the argument for the sake of completeness.

Proof. By the definition of E and the divergence theorem, we have that

$$\begin{aligned} 2(n-1)\omega_{n-1}E &= \lim_{r \rightarrow \infty} \int_{|x|=r} \sum_{i,j=1}^n (g_{ij,i} - g_{ii,j}) \nu_0^j \\ &= \int_{|x|=r} \sum_{i,j=1}^n (g_{ij,i} - g_{ii,j}) \nu_0^j \\ &\quad + \int_{|x| \geq r} \sum_{i,j=1}^n (g_{ij,ij} - g_{ii,jj}) \end{aligned}$$

for all r sufficiently large, and similarly for \bar{E} . All integrals here are with respect to the Euclidean metric. Note that $\sum_{i,j=1}^n (g_{ij,ij} - g_{ii,jj}) = 2\mu + O(|x|^{-2-2q})$, because both sides differ from the scalar curvature by terms quadratic in $\partial_k g_{ij}$ and π_{ij} . Since $(\mu - \bar{\mu}) \in L^1_{-n-q_0/2}$, it follows that there exists r_0 large and depending only on $\epsilon, n, p, q, q_0, \|\mu - \bar{\mu}\|_{L^1_{-n-q_0/2}}, \|(g - g_{\mathbb{E}}, \pi)\|_{W^{2,p}_{-q} \times W^{1,p}_{-q-1}}$, and $\|(\bar{g} - g_{\mathbb{E}}, \bar{\pi})\|_{W^{2,p}_{-q} \times W^{1,p}_{-q-1}}$ such that

$$\int_{|x| \geq r} \left| \sum_{i,j=1}^n (g_{ij,ij} - g_{ii,jj}) - (\bar{g}_{ij,ij} - \bar{g}_{ii,jj}) \right| < (n-1)\omega_{n-1}\epsilon$$

for all $r \geq r_0$. For the difference of the boundary integrals, note that

$$\begin{aligned} & \left| \int_{|x|=r} \sum_{i,j=1}^n [(g_{ij,i} - g_{ii,j}) - (\bar{g}_{ij,i} - \bar{g}_{ii,j})] \nu_0^j \right| \\ & \leq 2 \sum_{i,j,k=1}^n \int_{|x|=r} |\partial_k(g - \bar{g})_{ij}| \\ & \leq 2\omega_{n-1}r^{n-1} \sum_{i,j,k=1}^n \sup_{|x|=r} |\partial_k(g - \bar{g})_{ij}|(x) \\ & \leq 2(\omega_{n-1}r^{n-1})n^3(r^{-1-q}C_p\|g - \bar{g}\|_{W^{2,p}_{-q}}) \end{aligned} \tag{34}$$

where C_p is the constant that governs the continuous embedding $W^{1,p}_{-1-q} \subset C^{1-\frac{n}{p}}_{-1-q}$. By choosing $\delta > \|g - \bar{g}\|_{W^{2,p}_{-q}}$ sufficiently small (depending on r) we can ensure that (34) is less than $(n-1)\omega_{n-1}\epsilon$ so that $|E - \bar{E}| < \epsilon$. The argument for the linear momentum is similar. \square

6.1 Perturbing to strict dominant energy condition

We define the constraint map

$$\Phi(g, \pi) = (2\mu, J) = \left(R_g - |\pi|_g^2 + \frac{1}{n-1}(\text{tr}_g \pi)^2, \text{div}_g \pi \right).$$

We will also use the modified Lie derivative

$$\mathcal{L}_g Y = L_Y g - (\text{div}_g Y)g$$

of a vector field Y .

Lemma 20. *Let $(g - g_{\mathbb{E}}, \pi) \in W^{2,p}_{-q} \times W^{1,p}_{-1-q}$. The linear map $A : W^{2,p}_{-q} \times W^{1,p}_{-1-q} \rightarrow L^p_{-2-q}$ given by*

$$A(h, w) = D\Phi|_{(g, \pi)}(h, w) - (0, \frac{1}{2}h^{j\ell}J_\ell)$$

is surjective.

The proof is a small modification of the proof of surjectivity of the linearization $D\Phi|_{(g,\pi)}$ in [11, Proposition 3.1]. For the convenience of the reader, we include the argument here.

Proof. In this proof we treat Φ as a map defined on the space of (g, π) where g is a $(0, 2)$ -tensor and π is a $(2, 0)$ -tensor, and $(\operatorname{div}_g \pi)^i := (\pi^{ij})_{;j}$. By direct computation,

$$\begin{aligned} D\Phi|_{(g,\pi)}(h, w) = & \left(-\Delta_g(\operatorname{tr}_g h) + \operatorname{div}_g \operatorname{div}_g(h) - h^{ij} R_{ij} - 2h_{ij}\pi_\ell^i \pi^{j\ell} \right. \\ & - 2\pi_k^j w_j^k + \frac{2}{n-1} \operatorname{tr}_g \pi(h_{ij}\pi^{ij} + \operatorname{tr}_g w), \\ & \left. (\operatorname{div}_g w)^i - \frac{1}{2}\pi^{jk}h_{jk;\ell}g^{\ell i} + \pi^{jk}h_{j;k}^i + \frac{1}{2}\pi^{ij}(\operatorname{tr}_g h)_{,j} \right). \end{aligned}$$

Here all indices are raised or lowered with respect to g . The formula is well-known and can be found in, for example, [18, pp. 999–1000] for $n = 3$, but note that the negative divergence operator is used there. Let $(v, Z) \in W_{-q}^{2,p}$ where v is a function and Z is a vector field. Consider $h_{ij} = vg_{ij}$ and $w^{ij} = (\mathcal{L}_g Z)^{ij}$. Then

$$(v, Z) \mapsto A(h, w) \tag{35}$$

is a Fredholm operator from $W_{-q}^{2,p}$ to L_{-2-q}^p , cf. [5]. It follows that the range of A , which contains the range of the operator in (35), has finite codimension in L_{-2-q}^p . In particular, the range of A is closed.

Since A has closed range, we can prove surjectivity of A by showing that the kernel of the adjoint operator $A^*(\xi, Z) = D\Phi|_{(g,\pi)}^*(\xi, Z) - (\frac{1}{2}Z_i J_j, 0)$ is trivial. One can compute the formal L^2 -adjoint operator of $D\Phi|_{(g,\pi)}$

$$\begin{aligned} D\Phi|_{(g,\pi)}^*(\xi, Z) = & \left(-(\Delta_g \xi)g_{ij} + \xi_{;ij} - \xi R_{ij} + \left(\frac{2}{n-1}(\operatorname{tr}_g \pi)\pi_{ij} - 2\pi_{ik}\pi_j^k \right)\xi \right. \\ & + \frac{1}{2} \left((L_Z \pi)_{ij} + (Z_{;k}^k)\pi_{ij} - Z_i \pi_{j;k}^k - Z_j \pi_{i;k}^k - Z_{k;m}\pi^{km}g_{ij} - Z_k \pi_{;m}^{km}g_{ij} \right), \\ & \left. - \frac{1}{2}(L_Z g)^{ij} + \left(\frac{2}{n-1}(\operatorname{tr}_g \pi)g^{ij} - 2\pi^{ij} \right)\xi \right). \end{aligned}$$

Let (ξ, Z) be in the dual space $L_{-n+2+q}^{p^*}$ such that $A^*(\xi, Z) = (0, 0)$. Taking the trace of the first component of $A^*(\xi, Z) = (0, 0)$ gives an equation for $\Delta_g \xi$. Using this equation, we can eliminate the term $\Delta_g \xi$ from the system $A^*(\xi, Z) = (0, 0)$ to obtain

$$\begin{aligned} 0 = & \xi_{;ij} - \xi R_{ij} + \frac{1}{n-1}(R_g + 2|\pi|_g^2 - \frac{2}{n-1}(\operatorname{tr}_g \pi)^2)\xi g_{ij} \\ & + \left(\frac{2}{n-1}(\operatorname{tr}_g \pi)\pi_{ij} - 2\pi_{ik}\pi_j^k \right)\xi \\ & - \frac{1}{2(n-1)} \left(\operatorname{tr}_g(L_Z \pi) + Z_{;k}^k \operatorname{tr}_g \pi - 2Z^i \pi_{i;k}^k - Z_{k;m}\pi^{km} - Z_k \pi_{;m}^{km} \right) g_{ij} \\ & + \frac{1}{2} \left((L_Z \pi)_{ij} + (Z_{;k}^k)\pi_{ij} - Z_i \pi_{j;k}^k - Z_j \pi_{i;k}^k \right) \\ & - \frac{1}{2}Z_i J_j + \frac{1}{2(n-1)}Z^k J_k g_{ij} \\ 0 = & \frac{1}{2}(L_Z g)^{ij} + \left(2\pi^{ij} - \frac{2}{n-1}(\operatorname{tr}_g \pi)g^{ij} \right)\xi. \end{aligned} \tag{36}$$

We use a bootstrap argument to show that (ξ, Z) vanishes to infinite order at infinity, i.e. that $|(\xi, Z)| \leq C_N|x|^{-N}$ for any integer $N > 1$. The initial decay of ξ and Z is on the order of $|x|^{-n+2+q}$. By (36), $\nabla^2\xi$ is of order $|x|^{-n}$ and L_Zg is of order $|x|^{-n+1}$. The following estimates for $n = 3$ were proved in [11, (10)] and can be generalized easily to any $n \geq 3$. For any weight $\tau > 0$ and for radius $R > 0$ large,

$$\begin{aligned} \int_{M \setminus B_R} (\xi|x|^\tau)^2 |x|^{-n} &\leq C \int_{M \setminus B_R} (|\nabla^2\xi| |x|^{2+\tau})^2 |x|^{-n} \\ \int_{M \setminus B_R} (|Z||x|^\tau)^2 |x|^{-n} &\leq C \int_{M \setminus B_R} (|L_Zg| |x|^{1+\tau})^2 |x|^{-n}. \end{aligned}$$

By letting $\tau < n - 2$, we conclude that (ξ, Z) is of order $|x|^{-\tau}$ for any $\tau < n - 2$, so the decay rate of (ξ, Z) is improved. We can then argue inductively that (ξ, Z) vanishes to infinite order at infinity.

Finally, we take the trace of the first set of equations and the divergence of the second set of equations of (36). The leading order term in the divergence of the second equation is

$$(\operatorname{div}_g(L_Zg))^i = (Z_{;k}^i g^{kj} + Z_{;k}^j g^{ik})_{;j} = \Delta_g Z^i + g^{ik} Z_{;kj}^j.$$

We replace the term $g^{ik} Z_{;kj}^j$ by terms involving only $\nabla Z, \nabla \xi$, and ξ , using the trace of the second component $A^*(\xi, Z)$. Hence we obtain a system of linear equations of the form

$$\Delta_g(\xi, Z) = B(x)(\nabla\xi, \nabla Z) + C(x)(\xi, Z),$$

where B, C are coefficient matrices. Note that although $B(x)$ and $C(x)$ are different from the coefficient matrices in [11], they do have the same asymptotics. Using a Kelvin transform and a unique continuation argument as in [11, pp. 196–197], one sees that (ξ, Z) must vanish identically. This completes the proof of surjectivity. \square

Notation. For better readability, we define

$$s := \frac{4}{n-2},$$

which will appear frequently in this section.

For $(u-1, Y) \in W_{-q}^{2,p}$, let

$$\tilde{g} = u^s g \quad \text{and} \quad \tilde{\pi} = u^{s/2}(\pi + \mathcal{L}_g Y).$$

Define the operator

$$T(u, Y) = (2\tilde{\mu}u^s, \tilde{J}u^{s/2}),$$

where $(\tilde{\mu}, \tilde{J})$ are the mass and energy densities of $(\tilde{g}, \tilde{\pi})$. Explicitly, the components of T are given by

$$\begin{aligned} 2\tilde{\mu}u^s &= -\frac{4(n-1)}{n-2}u^{-1}\Delta_g u + R_g + \frac{1}{n-1}(\text{tr}_g \pi + \text{tr}_g \mathcal{L}_g Y)^2 \\ &\quad - (|\pi|_g^2 + 2(\mathcal{L}_g Y)_{kl}\pi^{kl} + |\mathcal{L}_g Y|_g^2) \\ \tilde{J}_j u^{s/2} &= (\text{div}_g \mathcal{L}_g Y + \text{div}_g \pi)_j + (n-1)\frac{s}{2}u^{-1}u_{,k}(\pi + \mathcal{L}_g Y)_j^k \\ &\quad - \frac{s}{2}u^{-1}u_{,j} \text{tr}_g(\pi + \mathcal{L}_g Y), \quad j = 1, 2, \dots, n. \end{aligned} \quad (37)$$

Here, all indices are raised and lowered with respect to g .

Lemma 21. *Suppose that $(g - g_{\mathbb{E}}, \pi) \in W_{-q}^{2,p} \times W_{-1-q}^{1,p}$. For any $(f, V) \in L_{-2-q}^p$ there exist $(v, Z) \in W_{-q}^{2,p}$ and symmetric $(0, 2)$ -tensors h, w in $C_c^{3,\alpha}$ so that*

$$DT|_{(1,0)}(v, Z) + D\Phi|_{(g,\pi)}(h, w) = (f, V_j + \frac{1}{2}h_j^\ell J_\ell).$$

If in addition $(f, V) \in C_{-n-q_0}^{0,\alpha}$ for some $q_0 > 0$ and $\alpha \in (0, 1 - \frac{n}{p}]$, and $(g, \pi) \in C_{loc}^{2,\alpha} \times C_{loc}^{1,\alpha}$, then we have $(v, Z) \in C_{2-n}^{2,\alpha}$.

Proof. The linearization of T at $(1, 0)$ is

$$\begin{aligned} DT|_{(1,0)}(v, Z) &= \left(-\frac{4(n-1)}{n-2}\Delta_g v - 4Z_{k;\ell}\pi^{k\ell} + \frac{2}{n-1}\text{tr}_g \pi \text{div}_g Z, \right. \\ &\quad \left. \text{div}_g(\mathcal{L}_g Z)_j + \frac{(n-1)s}{2}v_{,k}\pi_j^k - \frac{s}{2}v_{,j} \text{tr}_g \pi \right) \end{aligned} \quad (38)$$

where all covariant derivatives and raising of indices are with respect to g . For $0 < a < (n-2)$ and $p > n$, $DT|_{(1,0)} : W_{-a}^{2,p} \rightarrow L_{-2-a}^p$ is Fredholm with index zero, cf. [5]. Because the linear map A defined in Lemma 20 is surjective onto L_{-2-q}^p we can find $C^{3,\alpha}$ compactly supported symmetric tensor fields $\{(h_k, w_k)\}_{k=1}^N$ whose images $A(h_k, w_k)$ span a subspace that complements the image of $DT|_{(1,0)}$ in L_{-2-q}^p . It follows that for every $(f, V) \in L_{-2-q}^p$ there exist $(v, Z) \in W_{-q}^{2,p}$ and $(h, w) \in \text{span}\{(h_k, w_k)\}_{k=1}^N$ so that

$$DT|_{(1,0)}(v, Z) + D\Phi|_{(g,\pi)}(h, w) = (f, V_j + \frac{1}{2}h_j^\ell J_\ell). \quad (39)$$

This completes the first part of the proof.

Now suppose that $(f, V) \in C_{-n-q_0}^{0,\alpha}$ for some $q_0 > 0$, and $(g, \pi) \in C_{loc}^{2,\alpha} \times C_{loc}^{1,\alpha}$. We view the system (39) as $(n+1)$ Poisson equations in v, Z_i with a decaying nonhomogeneous term. Note that the contribution $D\Phi|_{(g,\pi)}(h, w)$ to the nonhomogeneous term is a compactly supported Hölder function. Using [24, Theorem 2], which treats decay of solutions of the Poisson equation, and combining it with weighted Schauder estimates and a bootstrapping argument, we can conclude that $(v, Z) \in C_{2-n}^{2,\alpha}$. \square

Theorem 22. *Let (M, g, π) be an asymptotically flat initial data set of type (p, q, q_0, α) . Assume that the dominant energy condition $\mu \geq |J|_g$ holds. For any $\delta > 0$, there exists asymptotically flat initial data $(\bar{g}, \bar{\pi})$ of the same type such that*

$$\|g - \bar{g}\|_{W_{-q}^{2,p}} \leq \delta \quad \|\pi - \bar{\pi}\|_{W_{-1-q}^{1,p}} \leq \delta,$$

and for some $\gamma > 0$ depending on δ ,

$$\bar{\mu} > (1 + \gamma)|\bar{J}|_{\bar{g}}.$$

Proof. Choose a smooth positive function f such that $f = |x|^{-n-\min(1,q_0)}$ near infinity, and let $(v, Z) \in C_{2-n}^{2,\alpha}$ and $(h, w) \in C_c^{3,\alpha}$ be a solution of the system

$$DT|_{(1,0)}(v, Z) + D\Phi|_{(g,\pi)}(h, w) = (2f, \frac{1}{2}h_j^l J_l),$$

whose existence is guaranteed by Lemma 21 (with $V \equiv 0$). Our goal is to show that for sufficiently small $t > 0$, the formula

$$\bar{g} = (1 + tv)^s(g + th) \quad \text{and} \quad \bar{\pi} = (1 + tv)^{s/2}(\pi + t\mathcal{L}_g Z + tw) \quad (40)$$

gives the desired initial data $(\bar{g}, \bar{\pi})$ in the statement of the theorem. Since we clearly have $\|g - \bar{g}\|_{W_{-q}^{2,p}} \leq \delta$ and $\|\pi - \bar{\pi}\|_{W_{-1-q}^{1,p}} \leq \delta$ for small enough t , it suffices to show that $\bar{\mu} > (1 + \gamma)|\bar{J}|_{\bar{g}}$ for some $\gamma > 0$ that depends on t .

In the following we denote $u = 1 + tv$ and define

$$\Phi_1(1 + tv, tZ, th, tw) = (2\bar{\mu}u^s, \bar{J}u^{s/2}).$$

By Taylor expansion,

$$\begin{aligned} \Phi_1(1 + tv, tZ, th, tw) \\ &= \Phi_1(1, 0, 0, 0) + tD\Phi_1|_{(1,0,0,0)}(v, Z, h, w) + \mathcal{R} \\ &= (2\mu, J) + tDT|_{(1,0)}(v, Z) + tD\Phi|_{(g,\pi)}(h, w) + \mathcal{R} \\ &= (2\mu, J) + t(2f, \frac{1}{2}h_i^k J_k) + \mathcal{R}, \end{aligned} \quad (41)$$

where there is a minor abuse of notation in the last line, and the remainder term $\mathcal{R} = \mathcal{R}(x, t)$ has the form

$$\mathcal{R}(x, t) = t \int_0^1 [D\Phi_1|_{(1,0,0,0)+rt(v,Z,h,w)} - D\Phi_1|_{(1,0,0,0)}] (v, Z, h, w) dr.$$

Claim. *We have that*

$$|\mathcal{R}(x, t)| \leq Ct^2(1 + |x|)^{2-2n} = O(t^2 f).$$

where C is a constant that does not depend on x or t .

Proof of claim. Clearly, $\mathcal{R}(x, t) = O(t^2)$, so it suffices to work outside a large ball B that contains the support of (h, w) . Note that when $x \in M \setminus B$ we have that $D\Phi_1|_{(1,0,0,0)+rt(v,Z,h,w)}(v, Z, h, w) = DT|_{(1,0)+rt(v,Z)}(v, Z)$ and hence

$$\mathcal{R}(x, t) = t \int_0^1 [DT|_{(1,0)+rt(v,Z)} - DT|_{(1,0)}] (v, Z) dr.$$

From (37) we see that the linearization of T at (u, Y) is given by

$$\begin{aligned} & DT|_{(u,Y)}(v, Z) \\ &= \left(\frac{4(n-1)}{n-2} (u^{-2} v \Delta_g u - u^{-1} \Delta_g v) \right. \\ &\quad + \frac{2}{n-1} (\text{tr}_g \pi + \text{tr}_g \mathcal{L}_g Y) \text{tr}_g \mathcal{L}_g Z - 2(\pi + \mathcal{L}_g Y)^{kl} (\mathcal{L}_g Z)_{kl}, \\ &\quad \text{div}_g (\mathcal{L}_g Z)_j + (n-1) \frac{s}{2} (vu^{-1})_{,k} (\pi + \mathcal{L}_g Y)_j^k + (n-1) \frac{s}{2} u^{-1} u_{,k} (\mathcal{L}_g Z)_j^k \\ &\quad \left. - \frac{s}{2} (vu^{-1})_{,j} \text{tr}_g (\pi + \mathcal{L}_g Y) - \frac{s}{2} u^{-1} u_{,j} \text{tr}_g \mathcal{L}_g Z \right). \end{aligned}$$

The claim follows from substituting $(1, 0) + rt(v, Z)$ and $(1, 0)$ for (u, Y) , using that $(v, Z) \in C_{2-n}^{2,\alpha}$, and estimating each term in an obvious way. \square

From (41) we have

$$u^s \bar{\mu} = \mu + tf + O(t^2 f) \quad \text{and} \quad u^{s/2} \bar{J}_i = J_i + t \frac{1}{2} h_i^k J_k + O(t^2 f).$$

In particular, for sufficiently small $t > 0$, we have

$$u^s \bar{\mu} > \mu + \frac{t}{2} f \quad \text{everywhere on } M. \tag{42}$$

We claim that $u^s |\bar{J}|_{\bar{g}} < |J|_g + \frac{tf}{4}$ for $t > 0$ sufficiently small. Choose f_1 to be a smooth nonnegative function whose support is larger than that of h . Then

$$\bar{g}^{ij} = u^{-s} (g^{ij} - th^{ij} + O(t^2 f_1)).$$

Therefore

$$\begin{aligned} (u^s |\bar{J}|_{\bar{g}})^2 &= u^{2s} \bar{g}^{ij} \bar{J}_i \bar{J}_j \\ &= (g^{ij} - th^{ij} + O(t^2 f_1)) (J_i + t \frac{1}{2} h_i^k J_k + O(t^2 f)) (J_j + t \frac{1}{2} h_j^k J_k + O(t^2 f)) \\ &= g^{ij} J_i J_j + t (-h^{ij} J_i J_j + \frac{1}{2} g^{ij} h_i^k J_k J_j + \frac{1}{2} g^{ij} J_i h_j^k J_k) \\ &\quad + O(t^2 |J|^2 f_1 + t^2 |J| f + t^3 f_1 + t^4 f^2) \\ &= |J|_g^2 + O(t^2 |J| f + t^3 f^2) \\ &= \left(|J|_g + \frac{tf}{4} \right)^2 - \frac{tf}{2} |J|_g - \frac{t^2 f^2}{16} + O(t^2 |J| f + t^3 f^2) \\ &< \left(|J|_g + \frac{tf}{4} \right)^2, \end{aligned}$$

where we choose $t > 0$ to be sufficiently small in the last line, proving the claim. Observe that the computation above explains the motivation behind the definition of A . The $\frac{1}{2}h_j^l J_l$ term in A is chosen specifically so that the first order change in $|J|_g$ under the deformation vanishes.

Now fix $t > 0$ small enough so that $u^s \bar{\mu} > \mu + \frac{tf}{2}$ and $u^s |\bar{J}|_{\bar{g}} < |J|_g + \frac{tf}{4}$. Our assumptions imply that $\sup_M \frac{|J|_g}{f} < \infty$. For $x \in M$ such that $|\bar{J}|_{\bar{g}}(x) \neq 0$ it follows that

$$\frac{\bar{\mu}}{|\bar{J}|_{\bar{g}}} = \frac{u^s \bar{\mu}}{u^s |\bar{J}|_{\bar{g}}} > \frac{\mu + tf/2}{|J|_g + tf/4} \geq \frac{|J|_g + tf/2}{|J|_g + tf/4} \geq 1 + \gamma$$

where $\gamma^{-1} := 1 + \frac{4}{t} \sup_M \frac{|J|_g}{f}$. Since $\bar{\mu} > 0$ we conclude that $\bar{\mu} > (1 + \gamma)|\bar{J}|_{\bar{g}}$ on M , as desired. Finally, note that γ depends only on t and $\sup_M \frac{|J|_g}{f}$. \square

Remark. The assertion of Lemma 1 in [32] is similar to but weaker than that of Theorem 22. We also note that the proof of this lemma contains an error.

6.2 Harmonic asymptotics

We first show that any asymptotically flat initial data set can be slightly perturbed so that

$$g = u^s g_{\mathbb{E}}, \quad \pi = u^{s/2} \mathcal{L}_{g_{\mathbb{E}}} Y$$

outside a compact set, for some choice of u and Y , meanwhile prescribing any constraints (μ, J) that are close to the original ones. We then show that (g, π) has harmonic asymptotics if the imposed constraints decay fast enough.

The first result is a modification of [11, Theorem 1], where the special case of three-dimensional vacuum initial data was treated.

Lemma 23. *Suppose $(g - g_{\mathbb{E}}, \pi) \in W_{-q}^{2,p} \times W_{-1-q}^{1,p}$. There exist $C_0, \delta_0 > 0$ so that given $(\bar{\mu}, \bar{J}) \in L_{-2-q}^p$ with $\|(\bar{\mu} - \mu, \bar{J} - J)\|_{L_{-2-q}^p} \leq \delta \leq \delta_0$, there is an initial data set $(\bar{g}, \bar{\pi})$ such that*

$$\bar{g} = u^s g_{\mathbb{E}}, \quad \bar{\pi} = u^{s/2} \mathcal{L}_{g_{\mathbb{E}}} Y \tag{43}$$

outside a compact set for some $(u-1, Y) \in W_{-q}^{2,p}$, and such that the mass and current densities of $(\bar{g}, \bar{\pi})$ are $(\bar{\mu}, \bar{J})$. Moreover,

$$\|\bar{g} - g\|_{W_{-q}^{2,p}} \leq C_0 \delta, \quad \|\bar{\pi} - \pi\|_{W_{-1-q}^{1,p}} \leq C_0 \delta.$$

Proof. For $\lambda \geq 1$ large define the cut-off initial data

$$\hat{g}_\lambda = \chi_\lambda g + (1 - \chi_\lambda) g_{\mathbb{E}}, \quad \hat{\pi}_\lambda = \chi_\lambda \pi$$

where $\chi_\lambda(x) = \chi(x/\lambda)$ and χ is a smooth cut-off function on \mathbb{R}^n that is 1 on $\{|x| \leq 1\}$ and 0 on $\{|x| \geq 2\}$. Note that $\|(\hat{g}_\lambda - g, \hat{\pi}_\lambda - \pi)\|_{W_{-q}^{2,p} \times W_{-1-q}^{1,p}} \rightarrow 0$ as $\lambda \rightarrow \infty$.

In the following, we suppress the subscript λ when the context is clear. Define

$$\tilde{g} = u^s \hat{g}, \quad \tilde{\pi} = u^{s/2} (\hat{\pi} + \mathcal{L}_{\hat{g}} Y),$$

where $(u-1, Y) \in W_{-q}^{2,p}$. Let $(\tilde{\mu}, \tilde{J})$ be the mass and current densities of $(\tilde{g}, \tilde{\pi})$. Define the map $T_{(\hat{g}, \hat{\pi})}(u, Y) = (2\tilde{\mu}, \tilde{J})$ from $(W_{-q}^{2,p} + 1) \times W_{-q}^{2,p} \rightarrow L_{-2-q}^p$. The linearization of $T_{(\hat{g}, \hat{\pi})}$ at $(1, 0)$ is

$$\begin{aligned} & DT_{(\hat{g}, \hat{\pi})}|_{(1,0)}(v, Z) \\ &= \left(-\frac{4(n-1)}{n-2} \Delta_{\hat{g}} v - s [R_{\hat{g}} - |\hat{\pi}|_{\hat{g}}^2 + \frac{1}{n-1} (\text{tr}_{\hat{g}} \hat{\pi})^2] v \right. \\ &\quad - 4Z_{k;l} \hat{\pi}^{kl} + \frac{2}{n-1} \text{tr}_{\hat{g}} \hat{\pi} \text{ div}_{\hat{g}} Z, \\ &\quad \left. \text{div}_{\hat{g}} (\mathcal{L}_{\hat{g}} Z)_j + (n-1) \frac{s}{2} v_{,k} \hat{\pi}_j^k - \frac{s}{2} v_{,j} \text{tr}_{\hat{g}} \hat{\pi} - \frac{s}{2} (\text{div}_{\hat{g}} \hat{\pi})_j v \right), \end{aligned}$$

where indices are raised and covariant derivatives are taken with respect to \hat{g} . The map $T_{(g, \pi)} : (W_{-q}^{2,p} + 1) \times W_{-q}^{2,p} \rightarrow L_{-2-q}^p$ is defined analogously. Because $q \in ((n-2)/2, n-2)$ and $p > n$, $DT_{(\hat{g}, \hat{\pi})}|_{(1,0)}$ and $DT_{(g, \pi)}|_{(1,0)}$ are Fredholm operators of index 0 for λ sufficiently large (see [5]).

Let K_1 be a complementing subspace for the kernel of $DT_{(g, \pi)}|_{(1,0)}$ in $W_{-q}^{2,p} \times W_{-2-q}^{1,p}$. Since the linearization $D\Phi|_{(g, \pi)} : W_{-q}^{2,p} \times W_{-1-q}^{1,p} \rightarrow L_{-2-q}^p$ is surjective (see [11] and Lemma 20) and because $DT_{(g, \pi)}|_{(1,0)}$ is Fredholm, we can find $C^{3,\alpha}$ compactly supported symmetric $(0, 2)$ -tensors $\{(h_k, w_k)\}_{k=1}^N$ whose images $D\Phi|_{(g, \pi)}(h_k, w_k)$ form a basis for a complementing subspace of the image of $DT_{(g, \pi)}|_{(1,0)}$ in L_{-2-q}^p . Let $K_2 = \text{span}\{(h_k, w_k)\}_{k=1}^N$. We define the maps $\bar{T}_{(\hat{g}, \hat{\pi})}, \bar{T}_{(g, \pi)} : K_1 \times K_2 \rightarrow L_{-2-q}^p$ by

$$\bar{T}_{(\hat{g}, \hat{\pi})}(u, Y, h, w) = \Phi(u^s \hat{g} + h, u^{s/2} (\hat{\pi} + \mathcal{L}_{\hat{g}} Y) + w)$$

and

$$\bar{T}_{(g, \pi)}(u, Y, h, w) = \Phi(u^s g + h, u^{s/2} (\pi + \mathcal{L}_g Y) + w).$$

The maps $\bar{T}_{(\hat{g}, \hat{\pi})}$ and $\bar{T}_{(g, \pi)}$ are continuously differentiable. Using that $(\hat{g}, \hat{\pi})$ converges to (g, π) in $W_{-q}^{2,p} \times W_{-q-1}^{1,p}$ as $\lambda \rightarrow \infty$ it is easy to see that $DT_{(\hat{g}, \hat{\pi})}|_{(u, Y, h, w)}$ converges to $DT_{(g, \pi)}|_{(u, Y, h, w)}$ as $\lambda \rightarrow \infty$ locally uniformly in $(u, Y, h, w) \in K_1 \times K_2$ in the strong operator topology. Observe that $DT_{(g, \pi)}|_{(1, 0, 0, 0)}$ is an isomorphism by construction. We conclude from the inverse function theorem that there exists $\delta_0 > 0$ such that for all $\lambda \geq 1$ sufficiently large, $\bar{T}_{(g, \pi)}$ restricts to a C^1 diffeomorphism defined on an open neighborhood of $(1, 0, 0, 0)$ (independent of $\lambda \geq 1$) in $K_1 \times K_2$ and onto an open neighborhood containing the L_{-2-q}^p ball of radius $2\delta_0$ centered at $(2\hat{\mu}, \hat{J})$.

Using that $\bar{T}_{(g,\pi)}(1, 0, 0, 0) = (2\mu, J)$ and that $\|(\mu, J) - (\hat{\mu}, \hat{J})\|_{L^p_{-2-q}} \rightarrow 0$ as $\lambda \rightarrow \infty$ we see that $\|(\bar{\mu}, \bar{J}) - (\hat{\mu}, \hat{J})\|_{L^p_{-2-q}} < 2\delta$ provided that $\|(\bar{\mu}, \bar{J}) - (\mu, J)\|_{L^p_{-2-q}} < \delta$ and $\lambda \geq 1$ is sufficiently large. Hence if $\delta \in (0, \delta_0)$ there exist $(u - 1, Y) \in W^{2,p}_{-q}$ and $C^{3,\alpha}$ compactly supported symmetric $(0, 2)$ -tensors (h, w) such that $\bar{T}_{(\hat{g}, \hat{\pi})}(u, Y, h, w) = (2\bar{\mu}, \bar{J})$ and such that

$$\|(u - 1, Y)\|_{W^{2,p}_{-q}} \leq C_1 \delta, \quad \|(h, w)\|_{W^{2,p}_{-q} \times W^{1,p}_{-1-q}} \leq C_1 \delta$$

for some constant $C_1 > 0$ that only depends on (g, π) . \square

The following proposition is straightforward except for a subtlety that arises when $n = 3$.

Proposition 24. *Suppose that (M, g, π) is an asymptotically flat initial data set of type (p, q, q_0, α) with $q_0 > 1$ (rather than just $q_0 > 0$) and such that*

$$g = u^s g_{\mathbb{E}}, \quad \pi = u^{s/2} \mathcal{L}_{g_{\mathbb{E}}} Y, \quad (44)$$

outside a large ball B for some $(u - 1, Y) \in W^{2,p}_{-q}$. Then (g, π) has harmonic asymptotics in the sense of Section 2.

Proof. By (37), outside a compact set, $(u - 1, Y)$ satisfies

$$\begin{aligned} \frac{4(n-1)}{n-2} u^{-1} \Delta_{g_{\mathbb{E}}} u - \frac{1}{n-1} (\text{tr}_{g_{\mathbb{E}}} \mathcal{L}_{g_{\mathbb{E}}} Y)^2 + |\mathcal{L}_{g_{\mathbb{E}}} Y|^2 &= -2\mu u^s \\ \Delta_{g_{\mathbb{E}}} Y_j + (n-1) \frac{s}{2} u^{-1} u_{,k} (\mathcal{L}_{g_{\mathbb{E}}} Y)_j^k - \frac{s}{2} u^{-1} u_{,j} \text{tr}_{g_{\mathbb{E}}} (\mathcal{L}_{g_{\mathbb{E}}} Y) &= J_j u^{s/2}, \end{aligned}$$

for $j = 1, 2, \dots, n$. By Sobolev embedding, $(u - 1, Y) = O^{1,\alpha}(|x|^{-q})$. As in the proof of Lemma 21, we may view this system as $n+1$ Poisson equations for the functions u, Y_i , so that $\Delta_{g_{\mathbb{E}}}(u, Y) = O^\alpha(|x|^{\max(-2q-2, -n-q_0)})$. Then using the fact that $q \in ((n-2)/2, n-2)$, Theorem 2 of [24] and weighted Schauder estimates imply that

$$\begin{aligned} u &= 1 + a|x|^{2-n} + O^{2,\alpha}(|x|^{2-n-\gamma}) \\ Y_i &= b_i |x|^{2-n} + O^{2,\alpha}(|x|^{2-n-\gamma}) \end{aligned} \quad (45)$$

for constants a, b_i , for any $0 < \gamma < \min(2q+2-n, q_0)$. It follows that $(u - 1, Y) = O^{2,\alpha}(|x|^{2-n})$.

For $n > 3$, since $\max(2-2n, -n-q_0) < -n-1$, we may repeat the above argument with improved decay of the source terms to conclude the desired result.

When $n = 3$, we argue as follows. We expand the nonlinear source terms above. We find that

$$\begin{aligned} |\mathcal{L}_{g_{\mathbb{E}}} Y|^2 &= |L_Y g|^2 - 2 \text{tr}_{g_{\mathbb{E}}} (L_g Y) (\text{div}_{g_{\mathbb{E}}} Y) + 3(\text{div}_{g_{\mathbb{E}}} Y)^2 \\ &= 2 \left[\sum_{i,j=1}^3 Y_{i,j}^2 + Y_{i,j} Y_{j,i} \right] - (\text{div}_{g_{\mathbb{E}}} Y)^2. \end{aligned}$$

Since

$$Y_{i,j} = -b_i \frac{x_j}{|x|^3} + O^{1,\alpha}(|x|^{-2-\gamma}),$$

we have

$$(\text{tr}_{g_{\mathbb{E}}} \mathcal{L}_{g_{\mathbb{E}}} Y)^2 = (\text{div}_{g_{\mathbb{E}}} Y)^2 = \left(\sum_{i=1}^3 Y_{i,i} \right)^2 = \frac{(b \cdot x)^2}{|x|^6} + O^{1,\alpha}(|x|^{-4-\gamma})$$

and

$$\sum_{i,j=1}^3 Y_{i,j}^2 + Y_{i,j} Y_{j,i} = \frac{|b|^2}{|x|^4} + \frac{(b \cdot x)^2}{|x|^6} + O^{1,\alpha}(|x|^{-4-\gamma}).$$

Thus the constraint equation for μ becomes

$$8\Delta_{g_{\mathbb{E}}} u = -\frac{2|b|^2}{|x|^4} - \frac{(b \cdot x)^2}{2|x|^6} + O^\alpha(|x|^{-4-\gamma}),$$

as long as $\gamma \leq q_0 - 1$. Now observe that

$$\begin{aligned} \Delta_{g_{\mathbb{E}}} \frac{x_i x_j}{|x|^4} &= \frac{2\delta_{ij}}{|x|^4} - \frac{4x_i x_j}{|x|^6}, \\ \Delta_{g_{\mathbb{E}}} \frac{1}{|x|^2} &= \frac{2}{|x|^4}. \end{aligned}$$

For appropriate choice of constants C_0 and C_{ij} , we have

$$\Delta_{g_{\mathbb{E}}} \left(u - C_0 \frac{1}{|x|^2} - C_{ij} \frac{x_i x_j}{|x|^4} \right) = O^\alpha(|x|^{-4-\gamma}).$$

Now apply [24, Theorem 2] and weighted Schauder estimates to obtain the desired result for u . The argument for Y is analogous. \square

6.3 Proof of Theorem 18

Using Theorem 22 and Proposition 19 we may reduce to the case where $\mu > (1 + \gamma)|J|_g$. For $l \geq 1$ define $\xi_\lambda(x) := \xi(\frac{x}{\lambda})$ where

$$\xi(x) = \begin{cases} 1 & \text{when } |x| \leq 1 \\ |x|^{-1} & \text{when } |x| \geq 2 \end{cases}.$$

Note that $\xi_\lambda(\mu, J) \in C_{-n-q_0-1}^{0,\alpha}$ and that $\|\xi_\lambda(\mu, J) - (\mu, J)\|_{L_{-2-q}^p}$ tends to zero as $\lambda \rightarrow \infty$. By Lemma 23 and Proposition 24 there are initial data (g_λ, π_λ) with harmonic asymptotics and mass and current densities $\xi_\lambda(\mu, J)$ such that $\|(g - g_\lambda, \pi - \pi_\lambda)\|_{W_{-q}^{2,p} \times W_{-1-q}^{1,p}} \rightarrow 0$ as $\lambda \rightarrow \infty$. In particular, $\|g - g_\lambda\|_{C^0} \rightarrow 0$ so that

$$|\xi_\lambda J|_{g_\lambda}^2 = g_\lambda^{ij} \xi_\lambda^2 J_i J_j = \xi_\lambda^2 |J|_g^2 (1 + o(1)) \quad \text{as } \lambda \rightarrow \infty.$$

It follows that

$$\xi_\lambda \mu > \xi_\lambda (1 + \gamma) |J|_g \geq \left(1 + \frac{\gamma}{2}\right) |\xi_\lambda J|_{g_\lambda}$$

for λ sufficiently large. In particular, the strict dominant energy condition holds for (g_λ, π_λ) . The convergence of mass and linear momentum as $\lambda \rightarrow \infty$ follows from Proposition 19. \square

Remark. It is a subtlety in the proof of Theorem 18 above that the decay conditions for (μ, J) improve when we are reducing to harmonic asymptotics. In fact, if we are content to preserve the dominant energy condition without requiring strictness, we can arrange for $(\bar{\mu}, \bar{J})$ to vanish outside a large compact set by choosing ξ with compact support. The modified initial data will then be smooth at infinity.

Remark. The assertion in [16] that $\text{tr}_g(k) = O(|x|^{-n})$ for n -dimensional initial data sets with harmonic asymptotics is wrong, as was pointed out to the first named author by A. Carlotto. When $n > 3$, the arguments in [16] continue to hold. When $n = 3$, the additional assumption that $\text{tr}_g(k) = O(|x|^{-\gamma})$ for some $\gamma > 2$ in Theorem 3 of [16] is required throughout (rather than just for the rigidity case). Of course, this is exactly the case treated by R. Schoen and S.-T. Yau in [32].

Remark. When $3 < n < 8$, the full positive mass theorem $E \geq |P|$ can be obtained from the positive energy theorem $E \geq 0$ in the form [16, Theorem 3] using the following reduction argument. Assume that $0 < E < |P|$. Fix $\theta \in (0, 1)$ such that $E' := \frac{E - \theta|P|}{1 - \theta^2} < 0$. Using the preceding remark we may assume that (M, g, k) is smooth (including decay on derivatives) outside a large compact subset of M and that $\mu = 0$ and $J = 0$ there. According to [7, Theorem 6.1], there exists a (vacuum) spacetime development of the asymptotically flat end of (M, g, k) in which we may deform the end of (M, g, k) to a boosted slice (of slope θ) that has energy E' . See [8, p. L115] for a remark on the transformation behavior of the energy-momentum tensor. The deformed initial data (M', g', k') satisfies the conditions of the positive energy theorem in [16, Theorem 3]. A contradiction.

We are grateful to P.T. Chruściel for useful discussions related to this argument. This reduction is folklore in the mathematical relativity community when the initial data set is given as a spacelike slice of an asymptotically flat spacetime, see e.g. [9].

Acknowledgements. The first named author acknowledges the support of NSF grant DMS-0906038 and of SNF grant 200021-140467. The second named author acknowledges the support of NSF grant DMS-1005560 and DMS-1308837. The third named author acknowledges the support of NSF grant DMS-0903467. The last named author acknowledges the support of NSF grant DMS-0604960.

References

- [1] William K. Allard, *On the first variation of a varifold: boundary behavior*, Ann. of Math. (2) **101** (1975), 418–446. MR 0397520 (53 #1379)
- [2] Lars Andersson, Marc Mars, and Walter Simon, *Local existence of dynamical and trapping horizons*, Phys. Rev. Lett. **95** (2005), 111102.
- [3] Lars Andersson and Jan Metzger, *The area of horizons and the trapped region*, Comm. Math. Phys. **290** (2009), no. 3, 941–972. MR 2525646 (2010f:53118)
- [4] Abhay Ashtekar and Gregory J. Galloway, *Some uniqueness results for dynamical horizons*, Adv. Theor. Math. Phys. **9** (2005), no. 1, 1–30. MR 2193368 (2006k:83101)
- [5] Robert Bartnik, *The mass of an asymptotically flat manifold*, Comm. Pure Appl. Math. **39** (1986), no. 5, 661–693. MR 849427 (88b:58144)
- [6] Robert Beig and Piotr T. Chruściel, *Killing vectors in asymptotically flat space-times. I. Asymptotically translational Killing vectors and the rigid positive energy theorem*, J. Math. Phys. **37** (1996), no. 4, 1939–1961. MR 1380882 (97d:83033)
- [7] D. Christodoulou and N. Ó Murchadha, *The boost problem in general relativity*, Comm. Math. Phys. **80** (1981), no. 2, 271–300. MR 623161 (84e:83011)
- [8] Piotr T. Chruściel, *A remark on the positive-energy theorem*, Classical Quantum Gravity **3** (1986), no. 6, L115–L121. MR 868711 (88f:83030a)
- [9] ———, *On the energy of the gravitational field at spatial infinity*, Conference on Mathematical Relativity (Canberra, 1988), Proc. Centre Math. Anal. Austral. Nat. Univ., vol. 19, Austral. Nat. Univ., Canberra, 1989, pp. 9–33. MR 1020788 (90k:83029)
- [10] Piotr T. Chruściel and Daniel Maerten, *Killing vectors in asymptotically flat space-times. II. Asymptotically translational Killing vectors and the rigid positive energy theorem in higher dimensions*, J. Math. Phys. **47** (2006), no. 2, 022502, 10. MR 2208148 (2007b:83054)
- [11] Justin Corvino and Richard M. Schoen, *On the asymptotics for the vacuum Einstein constraint equations*, J. Differential Geom. **73** (2006), no. 2, 185–217. MR 2225517 (2007e:58044)

- [12] Lu Ding, *Positive mass theorems for higher dimensional Lorentzian manifolds*, J. Math. Phys. **49** (2008), no. 2, 022504, 12. MR 2392853 (2008m:53081)
- [13] Frank Duzaar and Klaus Steffen, λ minimizing currents, Manuscripta Math. **80** (1993), no. 4, 403–447. MR 1243155 (95f:49062)
- [14] Michael Eichmair, *The Plateau problem for marginally outer trapped surfaces*, J. Differential Geom. **83** (2009), no. 3, 551–583. MR 2581357
- [15] ———, *Existence, regularity, and properties of generalized apparent horizons*, Comm. Math. Phys. **294** (2010), no. 3, 745–760. MR 2585986 (2011d:53171)
- [16] ———, *The Jang equation reduction of the spacetime positive energy theorem in dimensions less than eight*, Comm. Math. Phys. **319** (2013), no. 3, 575–593. MR 3040369
- [17] Michael Eichmair and Jan Metzger, *Jenkins-Serrin-type results for the Jang equation*, to appear in J. Differential Geom.
- [18] Arthur E. Fischer and Jerrold E. Marsden, *Linearization stability of the Einstein equations*, Bull. Amer. Math. Soc. **79** (1973), 997–1003. MR 0426035 (54 #13981)
- [19] Gregory J. Galloway, *Rigidity of marginally trapped surfaces and the topology of black holes*, Comm. Anal. Geom. **16** (2008), no. 1, 217–229. MR 2411473 (2009e:53087)
- [20] Gregory J. Galloway and Niall Ó Murchadha, *Some remarks on the size of bodies and black holes*, Classical Quantum Gravity **25** (2008), no. 10, 105009, 9. MR 2416045 (2009c:83058)
- [21] Gregory J. Galloway and Richard Schoen, *A generalization of Hawking’s black hole topology theorem to higher dimensions*, Comm. Math. Phys. **266** (2006), no. 2, 571–576. MR 2238889 (2007i:53078)
- [22] Lan-Hsuan Huang, *On the center of mass in general relativity*, Fifth International Congress of Chinese Mathematicians. Part 1, 2, AMS/IP Stud. Adv. Math., 51, pt. 1, vol. 2, Amer. Math. Soc., Providence, RI, 2012, pp. 575–591. MR 2908093
- [23] Joachim Lohkamp, *The higher dimensional positive mass theorem I*, arXiv:math/0608795v1 (2006).

- [24] Norman Meyers, *An expansion about infinity for solutions of linear elliptic equations*, J. Math. Mech. **12** (1963), 247–264. MR 0149072 (26 \#6568)
- [25] Thomas Parker and Clifford Henry Taubes, *On Witten’s proof of the positive energy theorem*, Comm. Math. Phys. **84** (1982), no. 2, 223–238. MR 661134 (83m:83020)
- [26] Richard Schoen, *Uniqueness, symmetry, and embeddedness of minimal surfaces*, J. Differential Geom. **18** (1983), no. 4, 791–809 (1984). MR 730928 (85f:53011)
- [27] ———, *Variational theory for the total scalar curvature functional for Riemannian metrics and related topics*, Topics in calculus of variations (Montecatini Terme, 1987), Lecture Notes in Math., vol. 1365, Springer, Berlin, 1989, pp. 120–154. MR 994021 (90g:58023)
- [28] ———, *Mean curvature in Riemannian geometry and general relativity*, Global theory of minimal surfaces, Clay Math. Proc., vol. 2, Amer. Math. Soc., Providence, RI, 2005, pp. 113–136. MR 2167257 (2006f:53044)
- [29] Richard Schoen and Leon Simon, *A new proof of the regularity theorem for rectifiable currents which minimize parametric elliptic functionals*, Indiana Univ. Math. J. **31** (1982), no. 3, 415–434. MR 652826 (84j:49039)
- [30] Richard Schoen and Shing Tung Yau, *On the proof of the positive mass conjecture in general relativity*, Comm. Math. Phys. **65** (1979), no. 1, 45–76. MR 526976 (80j:83024)
- [31] ———, *The energy and the linear momentum of space-times in general relativity*, Comm. Math. Phys. **79** (1981), no. 1, 47–51. MR 609227 (82j:83045)
- [32] ———, *Proof of the positive mass theorem. II*, Comm. Math. Phys. **79** (1981), no. 2, 231–260. MR 612249 (83i:83045)
- [33] Richard M. Schoen and Shing Tung Yau, *Complete manifolds with nonnegative scalar curvature and the positive action conjecture in general relativity*, Proc. Nat. Acad. Sci. U.S.A. **76** (1979), no. 3, 1024–1025. MR 524327 (80k:58034)
- [34] Leon Simon, *Lectures on geometric measure theory*, Proceedings of the Centre for Mathematical Analysis, Australian National University, vol. 3, Australian National University Centre for Mathematical Analysis, Canberra, 1983. MR 756417 (87a:49001)

- [35] Jean E. Taylor, *The structure of singularities in solutions to ellipsoidal variational problems with constraints in R^3* , Ann. of Math. (2) **103** (1976), no. 3, 541–546. MR 0428182 (55 #1208b)
- [36] Brian White, *The space of m -dimensional surfaces that are stationary for a parametric elliptic functional*, Indiana Univ. Math. J. **36** (1987), no. 3, 567–602. MR 905611 (88k:58027)
- [37] ———, *Existence of smooth embedded surfaces of prescribed genus that minimize parametric even elliptic functionals on 3-manifolds*, J. Differential Geom. **33** (1991), no. 2, 413–443. MR 1094464 (92e:58048)
- [38] Edward Witten, *A new proof of the positive energy theorem*, Comm. Math. Phys. **80** (1981), no. 3, 381–402. MR 626707 (83e:83035)